

ON THE POLYNOMIAL CONVERGENCE RATE TO NONEQUILIBRIUM STEADY-STATES

YAO LI

ABSTRACT. We consider a stochastic energy exchange model that models the 1D microscopic heat conduction in the nonequilibrium setting. In this paper, we prove the existence and uniqueness of the nonequilibrium steady state (NESS) and, furthermore, the polynomial speed of convergence to the NESS. Our result shows that the asymptotic properties of this model and its deterministic dynamical system origin are consistent. The proof uses a new technique called the induced chain method. We partition the state space and work on both the Markov chain induced by an “active set” and the tail of return time to this “active set”.

1. INTRODUCTION

As a ubiquitous process, heat conduction has been studied for over two hundred years. However, from a mathematical point of view, many microscopic aspects of heat conduction in solids are still unclear. For example, the derivation of macroscopic heat conduction laws like Fourier’s law from microscopic Hamiltonian dynamics is a well-known challenge in statistical mechanics for the past over a century. Over the last several decades, numerous mathematical models of 1-D microscopic heat conduction have been proposed and studied. Some of these models have purely deterministic dynamics [11, 18, 9, 32], while others are defined by stochastic differential equations [29, 24, 30, 31, 10, 2] or Markov jump processes [14, 20, 21, 33]. These models give mathematical frameworks for studying nonequilibrium phenomena including basic properties of nonequilibrium steady states (NESS), thermal conductivity, local thermodynamics equilibrium (LTE), fluctuation theorems, and eventually Fourier’s law.

This paper focuses on fundamental properties of nonequilibrium phenomena including the existence and uniqueness of the NESS and, furthermore, the polynomial-speed convergence to the NESS, for a class of 1-D microscopic heat conduction models. The model we study is a stochastic energy exchange model that is inspired by the KMP model introduced in [17], in which a chain of N sites are coupled with two heat baths. Each site carries a certain amount of energy. An exponential clock is associated with each pair of adjacent sites. When the clock rings, these two sites exchange energy in a “random halves” fashion. The energy exchange with the bath follows a similar rule. Different from the model in [17], the rate of an exponential clock here is energy-dependent. In this paper, the clock rate between two adjacent sites, called the *stochastic energy exchange rate*, depends on the square root of the minimum of two site energies. We refer Section 2.2 for the precise description of the model.




FIGURE 1. Locally confined particle system

The motivation of letting the stochastic energy exchange rate depend on the minimum of site energy comes from the study of a deterministic dynamical system heat conduction model, called the locally confined particle system. Introduced in [3], the locally confined particle system is a chain of locally confining cells in $\mathbb{R}^2$ like in Figure 1. An identical rigid disk-shaped moving particle is contained in each cell. A particle can not pass through the “bottleneck” between adjacent cells but can collide with its neighbors. Therefore, kinetic energy can be exchanged by particle-particle collisions. Studying such a purely deterministic dynamical system, especially in non-equilibrium settings, is very challenging. Only very limited rigorous results are known. On the other hand, it is well-known that chaotic billiard systems like the Lorentz gas have many stochastic properties due to the quick decay of correlation [4, 6, 5]. Hence a natural approach is to only record the kinetic energy of each particle and approximate this model by the stochastic energy exchange model described above. Let E_i and E_{i+1} be kinetic energies of adjacent particles. Assume the geometry of the cell allows a particle to be able to completely “hide” from its neighbors, i.e., neighboring particles can not collide with a particle when it is located at some area of its cell. Our numerical simulations in [19] show that the rate of interaction can be approximated by $\sim \sqrt{\min\{E_i, E_{i+1}\}}$ when one of E_i and E_{i+1} is sufficiently small. The heuristic reason of this rate is that when one particle is sufficiently slow and out the reach of its neighbors, the next particle-particle collision time should be primarily determined by the kinetic energy of the slow particle.

We remark that at a certain time rescaling limit, the locally confined particle system may have a different stochastic energy exchange rate. In a non-rigorous study of the locally confined particle system [13, 12], a rate function $\sim \sqrt{E_i + E_{i+1}}$ is obtained at a certain rare interaction limit and time rescaling limit. Assuming this rate of interaction, the mixing rate is known to be exponential [14, 20, 33]. Without any time rescaling limit, it is a simple mathematical fact that the speed of mixing in the locally confined particle system can not be faster than t^{-2} , provided particles can “hide” from their neighbors. This $\sim t^{-2}$ speed of mixing is also one of the main result of this paper. Therefore, the approximate interaction rate $\sim \sqrt{\min\{E_i, E_{i+1}\}}$ computed in [19] is consistent with the asymptotical dynamics of the locally confined particle system at its original time scale.

In this paper, among other results, we proved that when the stochastic rate of energy exchange between two sites are $\sim \sqrt{\min\{E_i, E_{i+1}\}}$, the Markov chain generated by our model has $\sim t^{-2}$ rate of mixing and $\sim t^{-2}$ rate of contraction. These

results completely match our analytical and numerical results about the locally confined particle system in [19]. As shown in the proof later in this paper, the main source of the slow-speed mixing comes from the rate $\sim \sqrt{\min\{E_i, E_{i+1}\}}$. When one site acquires a very low amount of energy from an energy exchange, the rates of two corresponding clocks become very low and can not be “rescued” by other “faster clocks”. Hence the next energy exchange at this site will not happen within a long time period, which obviously slows down the speed of mixing and convergence. We remark that this is also consistent with the mechanism of slow-speed mixing phenomenon of the locally confined particle system.

Despite the straightforward heuristic argument, a rigorous proof for the slow-speed mixing of a Markov process is known to be difficult. To the best of our knowledge, our result is the first polynomial convergence result in non-equilibrium settings. The closest related results we know are the slower-than-exponential convergence to the NESS in [35, 34, 7, 8] and the polynomial convergence to the equilibrium in [22]. In addition, we believe our results for the speed of mixing and contraction rate of the Markov operator are close to optimal. It is easy to see that the tail of the coupling time can not be less than t^{-2} , while the contraction rate is closely related to the coupling time. The speed of convergence to the NESS, which is shown to be faster than $\sim t^{-1}$, is not optimal due to the limitation of the proof. We will show that the speed of convergence to NESS is in fact also $\sim t^{-2}$ through numerical simulation in a separate paper.

The method of proving the polynomial-speed convergence to the steady-state is called the induced chain method. Since the source of slow convergence is the low-energy site, we partition the phase into two parts: the “active” set and the “inactive” set, where the “active” set means all site energies are above a certain threshold. Then we work on the Markov chain induced by the “active” set. Different from the model in [22], where the “active” set satisfies a Doeblin-type condition, in this paper we need some extra work to show the stochastic stability of the induced chain. The induced chain method consists of three steps. We first show that the induced chain admits a uniform reference set, on which trajectories can be coupled with strictly positive probability. Then we control the first passage time to this uniform reference set for the induced chain, this is done by constructing a Lyapunov function as we have done in [22]. Last, we show that the time duration of one step of the induced chain has a polynomial tail by constructing another Lyapunov function. The three steps above imply that the first passage time of the full system to the uniform reference set has a polynomial tail. Then we can apply results from discrete renewal theory and prove the polynomial-speed convergence and mixing.

We remark that a common way of proving polynomial-speed convergence is to construct a Lyapunov function [22, 15]. However, in this model such a construction is too complicated to be practical. One needs to consider both the lack of tightness and the possible inactive clocks in the construction of a Lyapunov function. By using the induced chain method, we can treat these two problems separately, and eventually give the tail of the first passage time to a uniform reference set. This method is useful in proving the polynomial (or sub-exponential) convergence of other

models. In addition, we believe the induced chain technique can be extended into a hierarchy of finitely many induced chains and be applied to a wider range of problems.

The paper is organized in the following way: Section 2 introduces the model and states the main result. The main strategy of proof, i.e., the induced chain method, is introduced in Section 3. Estimations for the time duration of one step of the induced chain are given in Section 4. The first entry time to the uniform reference set of the induced chain is done in Section 5. Finally, we complete the whole proof in Section 6.

2. MODEL AND RESULT

2.1. Stochastic approximation of deterministic dynamics. We start with a short review of the locally confined particle system and its stochastic approximation. Consider a chain of cells in $\mathbb{R}^2$ that are formed by finitely many piecewise C^3 curves. A rigid disk-shaped *moving particle* is confined in each cell, as shown in Figure 1. Two adjacent cells are connected by a “bottleneck” opening such that particles can not pass the opening but can collide with each other. A particle moves freely until it collides with the cell boundary or its neighbor particles. In addition, we assume each cell forms a strongly chaotic billiard table. In the absence of other particles, the billiard map of one particle is exponentially mixing. We refer to [3] for the ergodicity of the locally confined particle system under suitable conditions and [5] for major results of dynamic billiards.

In the locally confined particle system, a particle has a quick decay of correlation due to frequent collisions with the cell boundary. Therefore, it is natural to simplify the model by assuming that the geometry within a cell is forgotten by the particle. More precisely, we only record the kinetic energy carried by a particle and assume that the time to the next energy exchange is exponentially distributed. The rate of this exponential distribution is called the *stochastic energy exchange rate*, denoted by $R(E_i, E_{i+1})$, where E_i and E_{i+1} are kinetic energies of particles. If in addition, we assign a suitable rule for the energy redistribution in a particle-particle collision, a Markov jump process is obtained.

In [19], we numerically showed that the time to the next particle-particle collision always has an exponential tail that depends on the energy configuration of particles. This further supports the idea of approximating the locally confined particle system by a Markov jump process. The rate $R(E_i, E_{i+1})$ can be computed numerically as the slope of the exponential tail of the waiting time to the next particle-particle collision. Our numerical result in [19] showed that $R(E_i, E_{i+1}) \sim \sqrt{\min\{E_i, E_{i+1}\}}$ when at least one of E_i or E_{i+1} is sufficiently small.

As stated in the introduction, the slow convergence phenomenon comes from the slow clock rate when one of E_i or E_{i+1} is very small. To preserve this qualitative properties of the model, it is sufficient to let the stochastic energy exchange rate be $\sqrt{\min\{E_i, E_{i+1}\}}$ for all small $\min\{E_i, E_{i+1}\}$. Hence we assume $R(E_i, E_{i+1}) = \min\{K, \sqrt{\min(E_i, E_{i+1})}\}$ in this paper for the sake of simplicity. We idealize the energy exchange in a particle-particle collision by making it as a “random halves”

energy redistribution. In addition, the system is coupled with two heat baths and we prescribe a similar rule for the energy exchange with the heat bath. This gives rise to the stochastic energy exchange model as will be described in the next subsection.

2.2. Description of the stochastic energy exchange model. Now we will give a precise description of the stochastic energy exchange model. Consider a chain of N lattice sites connected to two heat baths at the ends. The energy at each site is denoted by $E_1, \dots, E_N$, respectively. The temperature of heat baths are T_L and T_R . An exponential clock is associated with each pair of adjacent sites. The rate of the clock depends on the energy at both sites, denoted by $R(E_i, E_{i+1})$. When the clock rings, the energy at two sites are pooled together and redistributed randomly as

$$(2.1) \quad (E'_i, E'_{i+1}) = (p(E_i + E_{i+1}), (1-p)(E_i + E_{i+1})),$$

where p is a uniform random variable distributed on $(0, 1)$ that is independent of everything else. In addition, an exponential clock is associated with the first (resp. last) site and the left (resp. right) heat bath, whose rate is $R(T_L, E_1)$ (resp. $R(E_N, T_R)$). When the clock rings, the energy at the first (resp. last) site exchanges energy with an exponential random variable :

$$E'_1 = p(E_1 + X_L) \quad (\text{resp. } E'_N = p(E_N + X_R)),$$

where p is a uniform random variable on $(0, 1)$ that is independent of everything else, X_L (resp. X_R) is an exponential random variable with mean T_L (resp. T_R). All exponential clocks are assumed to be mutually independent.

The rate $R(E_i, E_{i+1})$, called the *stochastic energy exchange rate* between sites i and $i + 1$, has the following form:

$$R(E_i, E_{i+1}) = \min\{K, \sqrt{\min(E_i, E_{i+1})}\},$$

where $K \gg 1$ is a sufficiently large constant. As explained above, the stochastic energy exchange rates is assumed to be the square root of the minimum of site energy when either of the site energy is sufficiently small. The maximum of stochastic energy exchange rate is set as $K < \infty$ for technical reasons. Without such an assumption, the Lyapunov function is not in the domain of the infinitesimal generator of the Markov chain, which impose certain technical complexity [25]. As the aim of this paper is to show that the property of the rate function at low energy leads to polynomial rate of convergence to NESS, we choose to cap the energy exchange rate by K . K is assumed to be sufficiently large so that it will not significant affect the dynamics at any “normal configuration”. In particular, we assume $K \gg T_L, T_R$.

It is easy to see from the description that this model generates a Markov jump process $\mathbf{E}_t = (E_1(t), \dots, E_N(t))$ on $\mathbb{R}_+^N$. The infinitesimal generator of $\mathbf{E}_t$ is

$$\begin{aligned} & \mathcal{L}f(E_1, \dots, E_N) \\ = & R(E_i, E_{i+1}) \sum_{i=1}^{N-1} \int_0^1 f(E_1, \dots, E_{i-1}, p(E_i + E_{i+1}), (1-p)(E_i + E_{i+1}), E_{i+2}, \dots, E_N) \\ & - f(E_1, \dots, E_N) dp \\ & + R(T_L, E_1) \int_0^\infty \int_0^1 \frac{1}{T_L} e^{-s/T_L} f(p(E_1 + s), E_2, \dots, E_N) - f(E_1, \dots, E_N) dp ds \\ & + R(E_N, T_R) \int_0^\infty \int_0^1 \frac{1}{T_R} e^{-s/T_R} f(E_1, \dots, E_{N-1}, p(E_N + s)) - f(E_1, \dots, E_N) dp ds \end{aligned}$$

We denote the transition kernel of $\mathbf{E}_t$ by $P^t(\mathbf{E}, \cdot)$, where $\mathbf{E} \in \mathbb{R}_+^N$. The left and right operator generated by P^t are

$$(P^t \eta)(\mathbf{E}) = \int_{\mathbb{R}_+^N} P^t(\mathbf{E}, d\mathbf{x}) \eta(\mathbf{x})$$

for a measurable function $\eta(\mathbf{E})$ on $\mathbb{R}_+^N$, and

$$(\mu P^t)(B) = \int_{\mathbb{R}_+^N} P^t(\mathbf{E}, B) \mu(d\mathbf{E})$$

for a probability measure μ on $\mathbb{R}_+^N$. We also use notations $\mathbb{P}_{\mathbf{E}}$ and $\mathbb{E}_{\mathbf{E}}$ for conditional probability and conditional expectation with respect to the initial condition $\mathbf{E}_0 = \mathbf{E}$.

In this paper we will use energy exchange events frequently. The event that the exponential clock between E_{i-1} and E_i rings at a time t (resp. at a stopping time τ) is denoted by $\mathcal{C}_i(t)$ (resp. $\mathcal{C}_i(\tau)$) for all $i = 1 \sim N+1$. In addition, for the sake of simplicity we denote $E_0 = T_L$ and $E_{N+1} = T_R$.

2.3. Main Result. To state our main result precisely, the following functions and measure classes are necessary. Let

$$W(\mathbf{E}) = \sum_{i=1}^N E_i.$$

For any $0 < \eta \ll 1$, let

$$V(\mathbf{E}) = V_\eta(\mathbf{E}) = \sum_{m=1}^N \sum_{i=1}^{N-m+1} \left(\sum_{j=0}^{m-1} E_{i+j} \right)^{a_m \eta - 1},$$

where $a_m = 1 - (2^{m-1} - 1)/(2^N - 1)$ for $m = 1 \sim N$. Let $\mathcal{M}_\eta$ be the collection of probability measure μ on $\mathbb{R}_+^N$ such that

$$\int_{\mathbb{R}_+^N} W(\mathbf{E}) + V(\mathbf{E}) \mu(d\mathbf{E}) < \infty.$$

$\mathcal{M}_\eta$ covers a large class of probability measures. For example, if $\mathbf{X}$ is a random energy configuration that has finite expectation and its density at the neighborhood

of the “boundary” $\{\mathbf{E} = (E_1, \dots, E_N) \mid E_i = 0 \text{ for some } i = 1 \sim N\}$ is uniformly bounded, then the probability measure induced by $\mathbf{X}$ belongs to $\mathcal{M}_\eta$ for any $\eta > 0$.

We have the following results regarding the stochastic stability of $\mathbf{E}_t$.

Theorem 1 (Polynomial contraction of Markov operator). *For any $\gamma > 0$, there exists $\eta > 0$ such that for any $\mu, \nu \in \mathcal{M}_\eta$,*

$$\lim_{t \rightarrow \infty} t^{2-\gamma} \|\mu P^t - \nu P^t\|_{TV} = 0,$$

where $\|\cdot\|_{TV}$ is the total variation norm.

Theorem 2 (Properties of the invariant measure). *There exists a unique invariant measure π that is absolutely continuous with respect to the Lebesgue measure. In addition, for any $\gamma > 0$, there exists $\eta > 0$ such that for any $\mu \in \mathcal{M}_\eta$,*

$$\lim_{t \rightarrow \infty} t^{1-\gamma} \|\mu P^t - \pi\|_{TV} = 0$$

A corollary of polynomial contraction of Markov operator is the rate of correlation decay.

Theorem 3 (Polynomial correlation decay). *For any $\gamma > 0$ there exists a $\eta > 0$ such that for any $\mu \in \mathcal{M}_\eta$, let functions ξ and $\zeta \in L^\infty(\mathbb{R}_+^N)$. Then*

$$\left| \int_{\mathbb{R}_+^N} (P^t \zeta)(\mathbf{E}) \xi(\mathbf{E}) \mu(d\mathbf{E}) - \int_{\mathbb{R}_+^N} (P^t \zeta)(\mathbf{E}) \mu(d\mathbf{E}) \int_{\mathbb{R}_+^N} \xi(\mathbf{E}) \mu(d\mathbf{E}) \right| = O(1) \cdot \left(\frac{1}{t^{2-\gamma}} \right)$$

as $t \rightarrow \infty$.

3. APPROACH TOWARDS POLYNOMIAL ERGODICITY

The purpose of this section is to introduce a general approach, called the induced chain method, towards the polynomial ergodicity of a Markov process. We introduce this method under the generic setting, as it can be potentially applied to other models. Throughout this section, we let Ψ_n be a discrete-time Markov chain on a measurable space $(X, \mathcal{B})$. The transition kernel of Ψ_n is $\mathcal{P}(x, \cdot)$.

For $A \in \mathcal{B}$, we let τ_A be the first passage time to A :

$$\tau_A = \inf\{n > 0 \mid \Psi_n \in A\}.$$

A set $A \in \mathcal{B}$ is said to be *accessible* if $\mathbb{P}_x[\tau_A < \infty] = 1$ for every $x \in X$.

We say a Markov chain is *irreducible* with respect to a measure ϕ on $\mathcal{B}$ if for any $A \in \mathcal{B}$ with $\phi(A) > 0$ and any $x \in X$, there exists n such that $\mathcal{P}^n(x, A) > 0$. In other word, every set with positive ϕ -measure is accessible. We refer readers to Chapter 4 of [26] for the well-definedness of irreducibility. In fact, if Ψ_n is irreducible with respect to some measure ϕ , then there exists a “maximal irreducible measure” ψ that is unique up to equivalence class.

3.1. Splitting, Coupling, and moments of return times. By the polynomial ergodicity of Ψ_n , we mean the polynomial rate of contraction of the Markov operator $\mathcal{P}$, the polynomial rate of convergence to the invariant measure of Ψ_n , and the polynomial rate of correlation decay of Ψ_n . To prove the polynomial ergodicity, the bound of return times to a certain *uniform reference set* $\mathfrak{C} \in \mathcal{B}$ is crucial.

Definition 3.1. A set $\mathfrak{C} \in \mathcal{B}$ is said to be a uniform reference set if it satisfies

$$\sup_{x \in \mathfrak{C}} \mathcal{P}(x, \cdot) \geq \eta \theta(\cdot),$$

where θ is a probability measure on $(X, \mathcal{B})$ and η is a strictly positive real number.

We call $\mathfrak{C}$ a uniform reference set because processes starting from $\mathfrak{C}$ have some uniform “common future”. If such a uniform reference set $\mathfrak{C}$ exists, Ψ_n can be converted to a new process $\tilde{\Psi}_n$ on a modified state space $\tilde{X} = X \cup \mathfrak{C}_1$, where $\mathfrak{C}_1$ is an identical copy of $\mathfrak{C}_0 := \mathfrak{C}$. $\mathcal{B}$ can be extended to $\tilde{\mathcal{B}}$ on $\tilde{X}$ accordingly. Then we can split a probability measure μ on $(X, \mathcal{B})$ to a probability measure μ^* on $(\tilde{X}, \tilde{\mathcal{B}})$:

$$\begin{cases} \mu^*|_X = (1 - \eta) \mu|_{\mathfrak{C}_0} + \mu|_{X \setminus \mathfrak{C}_0} \\ \mu^*|_{\mathfrak{C}_1} = \eta \mu|_{\mathfrak{C}_0}, \quad \mathfrak{C}_0 \cong \mathfrak{C}_1 \text{ via the natural identification.} \end{cases}$$

Then we can “lift” Ψ_n to a new Markov process $\tilde{\Psi}_n$ on $\tilde{X}$ with a transition kernel $\tilde{\mathcal{P}}(x, \cdot)$:

$$\begin{cases} \tilde{\mathcal{P}}(x, \cdot) = (\mathcal{P}(x, \cdot))^* & x \in X \setminus \mathfrak{C}_0 \\ \tilde{\mathcal{P}}(x, \cdot) = [(\mathcal{P}(x, \cdot))^* - \eta \theta^*(\cdot)] / (1 - \eta) & x \in \mathfrak{C}_0 \\ \tilde{\mathcal{P}}(x, \cdot) = \theta^*(\cdot) & x \in \mathfrak{C}_1 \end{cases}$$

It is straightforward to check that $\tilde{\Psi}_n$ has an atom $\alpha := \mathfrak{C}_1$, i.e., $\tilde{\mathcal{P}}(x, \cdot)$ is the same for all $x \in \alpha$. In addition, $\tilde{\Psi}_n$ projects to Ψ_n through the natural projection from $\tilde{X}$ to X . This transformation is called the Nummelin splitting [27]. We refer [27, 26] for the detail.

If Ψ_n admits an accessible uniform reference set, many results about the stochastic stability of Ψ_n can be implied by estimates about $\tau_{\mathfrak{C}}$. To state the results, we need the aperiodicity of Ψ_n .

Definition 3.2. An irreducible Markov process Ψ_n is said to be *strongly aperiodic* if Ψ_n admits a uniform reference set $\mathfrak{C}$ such that $\theta(\mathfrak{C}) > 0$.

If Ψ_n is strongly aperiodic, Ψ_n is aperiodic such that no cyclic decomposition is possible. We refer Chapter 5 of [26] for the precise definition of aperiodicity of Markov processes on measurable state spaces.

Theorem 3.3 (Theorem 10.0.1 in [26]). *Let Ψ_n be an irreducible aperiodic Markov chain on $(X, \mathcal{B})$ with transition kernel $\mathcal{P}$. If $\mathfrak{C} \in \mathcal{B}$ is an accessible uniform reference set such that*

$$\sup_{x \in \mathfrak{C}} \mathbb{E}_x[\tau_{\mathfrak{C}}] < \infty,$$

then there exists an invariant probability measure π .

Beyond the existence, the polynomial ergodicity of Ψ_n follows from the finiteness of moments of $\tau_{\mathfrak{C}}$.

Theorem 3.4 (Theorem 2.7 in [28]). *Let Ψ_n be an aperiodic Markov chain on $(X, \mathcal{B})$ with transition kernel $\mathcal{P}$. Let π be an invariant probability measure of Ψ_n . If $\mathfrak{C} \in \mathcal{B}$ is an accessible uniform reference set and*

$$\sup_{x \in \mathfrak{C}} \mathbb{E}_x[\tau_{\mathfrak{C}}^\beta] < \infty$$

for some $\beta > 0$, then for any probability measures μ, ν on X that satisfy

$$\mathbb{E}_\mu[\tau_{\mathfrak{C}}^\beta] < \infty \quad \text{and} \quad \mathbb{E}_\nu[\tau_{\mathfrak{C}}^\beta] < \infty,$$

we have

$$\lim_{n \rightarrow \infty} n^\beta \|\mu \mathcal{P}^n - \nu \mathcal{P}^n\|_{TV} = 0.$$

In addition, if $\beta > 1$, then

$$\lim_{n \rightarrow \infty} n^{\beta-1} \|\mu \mathcal{P}^n - \pi\|_{TV} = 0$$

for any μ satisfies

$$\mathbb{E}_\mu[\tau_{\mathfrak{C}}^\beta] < \infty.$$

We give a short proof for Theorem 3.4 based on the discrete renewal theory in [23]. We refer [28, 22] for the full details, and [15] for a modern treatment of continuous-time Feller processes.

Proof. We first apply the Nummerlin splitting to Ψ_n to obtain $\tilde{\Psi}_n$ on $\tilde{X}$. $\tilde{\Psi}_n$ possesses an accessible atom α .

Let $\tilde{\Psi}_n^1$ and $\tilde{\Psi}_n^2$ be two independent copies of $\tilde{\Psi}_n$ starting from μ^* and ν^* , respectively. Let $Y_0^1, Y_1^1, Y_2^1, \dots$ and $Y_0^2, Y_1^2, Y_2^2, \dots$ be the passage times to α for two independent processes, respectively. Since α is an atom, $\{Y_i^1\}_{i=1}^\infty$ and $\{Y_i^2\}_{i=1}^\infty$ are i.i.d random variables. Therefore

$$S_n^1 := \sum_{i=0}^n Y_i^1, \text{ and } S_n^2 := \sum_{i=0}^n Y_i^2$$

form two delayed renewal processes. Y_0^1 and Y_0^2 are called the *delay distributions* of the renewal processes. Let T be the first simultaneous renewal time

$$T = \inf_{m \geq 0} \{S_{k_1}^1 = S_{k_2}^2 = m \text{ for some } k_1, k_2\}, .$$

Since Ψ_n (and $\tilde{\Psi}_n$) is aperiodic, renewal processes S_n^1 and S_n^2 are aperiodic.

Then it is easy to see that after T , $\tilde{\Psi}_n^1$ and $\tilde{\Psi}_n^2$ become indistinguishable. T is called the *coupling time* of $\tilde{\Psi}_n^1$ and $\tilde{\Psi}_n^2$. It is well-known that

$$\|\mu \mathcal{P}^n - \nu \mathcal{P}^n\|_{TV} \leq \|\mu^* \tilde{\mathcal{P}}^n - \nu^* \tilde{\mathcal{P}}^n\|_{TV} \leq 2\mathbb{P}_{\mu, \nu}[T > n].$$

Therefore it is sufficient to show the polynomial tail of $\mathbb{P}[T > n]$. Instead of the polynomial tail, we first prove the finiteness of moments of T by using the delayed renewal processes constructed above and the following two lemmas.

Lemma 3.5 (Lemma 3.1 of [28]). *Let Ψ_n be an aperiodic Markov chain on $(X, \mathcal{B})$ with transition kernel $\mathcal{P}$. If $\mathfrak{C} \in \mathcal{B}$ is an accessible uniform reference set and*

$$\sup_{x \in \mathfrak{C}} \mathbb{E}_x[\tau_{\mathfrak{C}}^\beta] < \infty$$

for some $\beta > 0$, then for any probability measure μ such that

$$\mathbb{E}_\mu[\tau_{\mathfrak{C}}^\beta] < \infty,$$

there exists a constant $C < \infty$, such that

$$\mathbb{E}_{\mu^*}[\tau_\alpha^\beta] \leq C \mathbb{E}_\mu[\tau_{\mathfrak{C}}^\beta] < \infty.$$

Lemma 3.6 (Section II.4 and II.5 of [23]). *Let S_n^1 and S_n^2 be the delayed renewal processes as above. If there exists $\beta > 1$ such that $\mathbb{E}[(Y_0^1)^\beta]$, $\mathbb{E}[(Y_0^2)^\beta]$, and $\mathbb{E}[(Y_1^1)^\beta]$ are all finite, then there exists a constant $C < \infty$, such that*

$$\mathbb{E}[T^\beta] < C(\mathbb{E}[(Y_0^1)^\beta] + \mathbb{E}[(Y_0^2)^\beta]) < \infty.$$

In addition, there exists a delay distribution Y_0^c with $\mathbb{E}[(Y_0^c)^{\beta-1}] < \infty$ such that

$$S_n^c := Y_0^c + \sum_{i=1}^n Y_i^c, \quad n \geq 1$$

is a stationary renewal process, where $\{Y_i^c\}_{i=1}^n$ are independent random variables that have the same distribution as Y_1^1 .

Bounds of $\mathbb{E}[(Y_0^1)^\beta]$, $\mathbb{E}[(Y_0^2)^\beta]$, and $\mathbb{E}[(Y_1^1)^\beta]$, i.e., $\mathbb{E}_{\mu^*}[\tau_\alpha^\beta]$, $\mathbb{E}_{\nu^*}[\tau_\alpha^\beta]$, and $\mathbb{E}_\alpha[\tau_\alpha^\beta]$, are given in Lemma 3.5. Bounds of $\mathbb{E}[T^\beta]$ when starting from μ and ν follow from Lemma 3.6. When starting from π , passage times to α form a (delayed) stationary renewal process. The delay distribution of the corresponding renewal process must be Y_0^c , as the delay distribution that lead to a stationary renewal process is unique ([1], Chapter 2). Therefore, bounds of $\mathbb{E}[T^\beta]$ when starting from μ and π also follow from Lemma 3.6.

Theorem 3.4 is then implied by the following simple probability fact. (See Lemma 3.9). Let Z be any nonnegative integer-valued random variable. For any $\beta > 0$,

$$\mathbb{E}[Z^\beta] < \infty \Rightarrow \lim_{n \rightarrow \infty} n^\beta \mathbb{P}[Z > n] = 0.$$

□

We remark that the bounds in Lemma 3.5 and Lemma 3.6 are more precise than those in the original statement. These bounds are used to prove the following corollary that is used in the proof of Theorem 3.

Corollary 3.7. *Let Ψ_n and $\mathfrak{C}$ be as in Theorem 3.4, then for any probability measures μ, ν on X that satisfy*

$$\mathbb{E}_\mu[\tau_{\mathfrak{C}}^\beta] < \infty \quad \text{and} \quad \mathbb{E}_\nu[\tau_{\mathfrak{C}}^\beta] < \infty,$$

there exists a constant C such that

$$\sup_n n^\beta \|\mu \mathcal{P}^n - \nu \mathcal{P}^n\|_{TV} \leq C(\mathbb{E}_\mu[\tau_{\mathfrak{C}}^\beta] + \mathbb{E}_\nu[\tau_{\mathfrak{C}}^\beta]),$$

Proof. We have

$$\sup_n n^\beta \|\mu \mathcal{P}^n - \nu \mathcal{P}^n\|_{TV} \leq 2 \sup_n n^\beta \mathbb{P}[T > n] \leq 2 \mathbb{E}_{\mu^*, \nu^*}[T^\beta].$$

The corollary then follows immediately from Lemma 3.5 and Lemma 3.6. $\square$

3.2. Induced chain method. As discussed above, a crucial step towards polynomial ergodicity is to estimate the moment of $\tau_{\mathfrak{C}}$, i.e., the first passage time to a certain uniform reference set. In some simple cases such as the model in [22], this can be done by constructing a Lyapunov function. However, in our model and many other problems, it is extremely difficult to find a single Lyapunov function to complete this task. Here, we introduce a method, called the induced chain method, that can be used to estimate the moment of $\tau_{\mathfrak{C}}$ under more general settings.

Let $X = G \cup B$ be a partition of the state space of Ψ_n , where G is the “good set” on which Ψ_n is sufficiently “active”, while B is the “bad set” on which Ψ_n may hover for a long time. Define $0 = T_0 < T_1 < \dots < T_n < \dots$ be return times to G such that

$$T_{n+1} = \inf_{k > T_n} \{\Psi_k \in G\}.$$

and let $\hat{\Psi}_n = \Psi_{T_n}$. Then it is easy to check that $\hat{\Psi}_n$ is a Markov chain induced by G .

Assume $\mathfrak{C} \subset G$. The tail of $\tau_{\mathfrak{C}}$ can be estimated by the following two assumptions.

- (i) $T_{n+1} - T_n$ has a polynomial tail for each T_n . There exists a constant $\alpha > 0$ such that

$$\mathbb{P}[T_{n+1} - T_n > k \mid \Psi_{T_n}] \leq \xi(\Psi_{T_n}) k^{-\alpha},$$

where $1 \leq \xi(\Psi_{T_n}) < \infty$ is a constant depending on Ψ_{T_n} . Furthermore, $\xi(\Psi_{T_n})$ is uniformly bounded by $\xi_1 < \infty$ for $\Psi_{T_n} \in G$.

- (ii) $\hat{\tau}_{\mathfrak{C}} = \inf_{n > 0} \{\hat{\Psi}_n \in \mathfrak{C}\}$ has an exponential tail. There exist constants $\omega, \eta = \eta(\Psi_0) > 0$ such that

$$\mathbb{P}_{\Psi_0}[\hat{\tau}_{\mathfrak{C}} > k] \leq \eta(\Psi_0) e^{-\omega k},$$

where $\eta(\Psi_0)$ depends on Ψ_0 .

Theorem 3.8. *Assuming (i) and (ii) above, then for any $\epsilon > 0$, there exists a constant c such that*

$$\mathbb{P}_{\Psi_0}[\tau_{\mathfrak{C}} > n] \leq c(\eta(\Psi_0) + \xi(\Psi_0)) n^{-(\alpha - \epsilon)}$$

for any $\Psi_0 \in X$.

Proof. For any small $\delta > 0$, we have

$$\{\tau_{\mathfrak{C}} > k^{1+\delta}\} \subset \{\hat{\tau}_{\mathfrak{C}} > k^\delta\} \bigcup_{n=0}^{\lfloor k^\delta \rfloor} \{T_{n+1} - T_n > k, \hat{\tau}_{\mathfrak{C}} > n\}.$$

This implies

$$\begin{aligned} \mathbb{P}_{\Psi_0}[\tau_{\mathfrak{C}} > k^{1+\delta}] &\leq \mathbb{P}_{\Psi_0}[\hat{\tau}_{\mathfrak{C}} > k^\delta] \\ &\quad + \sum_{n=0}^{\lfloor k^\delta \rfloor} \mathbb{P}_{\Psi_0}[T_{n+1} - T_n > k \mid \hat{\tau}_{\mathfrak{C}} > n]. \end{aligned}$$

By assumption (i) and the Markov property, we have

$$\mathbb{P}_{\Psi_0}[T_1 - T_0 > k \mid \hat{\tau}_{\mathfrak{C}} > 0] \leq \xi(\Psi_0)k^{-\alpha}$$

for $n = 0$ and

$$\begin{aligned} &\mathbb{P}_{\Psi_0}[T_{n+1} - T_n > k \mid \hat{\tau}_{\mathfrak{C}} > n] \\ &= \int_G \mathbb{P}[T_{n+1} - T_n > k \mid \Psi_{T_n} = x, \hat{\tau}_{\mathfrak{C}} > n] \mathbb{P}_{\Psi_0}[\Psi_{T_n} = dx \mid \hat{\tau}_{\mathfrak{C}} > n] \\ &\leq \xi_1 n^{-\alpha} \end{aligned}$$

for $n \geq 1$.

By assumption (ii), we have

$$\mathbb{P}_{\Psi_0}[\hat{\tau}_{\mathfrak{C}} > k^\delta] \leq \eta(\Psi_0)e^{-\omega k^\delta}.$$

Therefore,

$$\begin{aligned} &\mathbb{P}_{\Psi_0}[\tau_{\mathfrak{C}} > k^{1+\delta}] \\ &\leq \eta(\Psi_0)e^{-\omega k^\delta} + \max\{\xi(\Psi_0), \xi_1\}k^\delta k^{-\alpha} \\ &\leq c(\delta)(\eta(\Psi_0) + \xi(\Psi_0))n^{-(\alpha-\delta)} \end{aligned}$$

for some $c(\delta) > 0$ that depends on δ , as $e^{-\omega k^\delta}$ decays faster than $k^{-\alpha}$ when $k \rightarrow \infty$.

The theorem then follows by making $\delta > 0$ sufficiently small and let $c = c(\delta)$ for the δ we choose. $\square$

The finiteness of moments of a random variable is closely related to its polynomial tails. We finish this subsection by proving two simple probabilistic facts that will be used frequently in this paper.

Let Z be a random variable that takes nonnegative integer values in, and let $\beta > 1$.

Lemma 3.9. $\mathbb{E}[Z^\beta] < \infty \implies \lim_{n \rightarrow \infty} n^\beta \mathbb{P}[Z > n] = 0$.

Proof. First,

$$\begin{aligned} \sum_{n=0}^{\infty} n^{\beta-1} \mathbb{P}[Z > n] &= \sum_{n=0}^{\infty} n^{\beta-1} \sum_{m=n+1}^{\infty} \mathbb{P}[Z = m] \\ &= \sum_{m=1}^{\infty} \left(\sum_{n=0}^{m-1} n^{\beta-1} \right) \mathbb{P}[Z = m] \leq \text{constant} \cdot \mathbb{E}[Z^\beta]. \end{aligned}$$

Then, letting $a_n = \mathbb{P}[Z > n]$ so that $a_0 \geq a_1 \geq \dots$, we claim that

$$\text{if } \sum_{n=0}^{\infty} a_n n^{\beta-1} < \infty \quad \text{then} \quad a_n n^{\beta} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

To see that, write

$$\sum_{n=0}^{\infty} (n+1)^{\beta} (a_n - a_{n+1}) = \sum_{n=0}^{\infty} [(n+1)^{\beta} - n^{\beta}] a_n,$$

so our hypothesis implies the sum on the left side converges. Since

$$\sum_{n \geq N} (n+1)^{\beta} (a_n - a_{n+1}) = a_N (N+1)^{\beta} + \sum_{n \geq N+1} a_n ((n+1)^{\beta} - n^{\beta}),$$

it follows that both terms on the right tend to 0 as $N \rightarrow \infty$. $\square$

Lemma 3.10. *If $\mathbb{P}[Z > n] \leq C(n+1)^{-\beta}$ for $\beta > 1$, then for any $\varepsilon > 0$, there exists a constant K that depends on β and ε , such that $\mathbb{E}[Z^{\beta-\varepsilon}] < KC$.*

Proof. There exists a constant K_0 that only depends on ε and β , such that

$$\begin{aligned} \mathbb{E}[Z^{\beta-\varepsilon}] &= \sum_{n=0}^{\infty} n^{\beta-\varepsilon} \mathbb{P}[Z = n] \\ &\leq K_0 \sum_{n=0}^{\infty} \sum_{m=0}^n m^{\beta-1-\varepsilon} \mathbb{P}[Z = n] \\ &= K_0 \sum_{n=0}^{\infty} (n+1)^{\beta-1-\varepsilon} \mathbb{P}[Z > n] \\ &\leq K_0 \sum_{n=0}^{\infty} (n+1)^{\beta-1-\varepsilon} C(n+1)^{-\beta}. \end{aligned}$$

Hence there exists a constant K that depends on β and ε , such that the last quantity is less than KC . $\square$

We remark the induced chain method can be extended to a hierarchical setting. Let $X = B \cup G$ be the same partition as above. If we have $\mathfrak{C} := G_m \subset G_{m-1} \subset \dots \subset G_0 := G$ for $m \geq 1$, and the first return time to G_{i+1} for the G_i -induced chain has exponential tail for each $i = 0 \sim m-1$, then the similar argument in Theorem 3.8 still follows.

3.3. Lyapunov function and moments of return time. In the induced chain argument above, it remains to find sufficient conditions to estimate tails of $\hat{\tau}_{\mathfrak{C}}$ and $T_{n+1} - T_n$. This can be done by constructing Lyapunov-type functions. We introduce the following two theorems that will be used later.

Theorem 3.11. *(Theorem 15.2.5 of [26])*

Let Ψ_n be a Markov chain on $(X, \mathcal{B})$ with transition kernel $\mathcal{P}$. We assume that there exist a function $W : X \rightarrow [1, \infty]$, a set $A \in \mathcal{B}$, constants $b, c > 0$ and $0 \leq \beta < 1$ such that

$$\mathcal{P}W - W \leq -\beta W + b\mathbf{1}_A.$$

then for any $r \in (1, (1 - \beta)^{-1})$, there exists $\epsilon > 0$ such that

$$\mathbb{E}_x \left[\sum_{k=0}^{\tau_A-1} W(\Psi_k) r^k \right] \leq \epsilon^{-1} r^{-1} W(x) + \epsilon^{-1} b \mathbf{1}_A.$$

Theorem 3.12. (Theorem 3.6 of [16].) Let Ψ_n be a Markov chain on $(X, \mathcal{B})$ with transition kernel $\mathcal{P}$. We assume that there exist a function $W : X \rightarrow [1, \infty]$, a set $A \in \mathcal{B}$, constant $b > 0$ and $0 \leq \beta < 1$ such that

$$\mathcal{P}W - W \leq -cW^\beta + b\mathbf{1}_A.$$

Then there is a constant $\hat{c}$ such that for all $x \in X$,

$$\mathbb{E}_x \left[\sum_{k=0}^{\tau_A-1} (k+1)^{\hat{\beta}-1} \right] \leq \hat{c} W(x), \quad \hat{\beta} = (1 - \beta)^{-1}.$$

The reader may notice that some assumptions in the original theorems in [16] and [26] are skipped in Theorem 3.11 and Theorem 3.12. This is because we only need the first time to the “bottom” of the Lyapunov function, while both [16] and [26] also consider first passage times to *any* set.

4. EXCURSION TIME ON LOW ENERGY SET

It is obvious that for $\mathbf{E}_t$, the “bad set” B (see Section 3.2) should consist of energy configurations at which at least one of E_i is sufficiently small. It remains to estimate the excursion time on this “bad set”. To do so, we define the following sequence and function. Let $0 < \eta \ll 1$ be a parameter that will be determined later. Let $a_1, \dots, a_N$ be the sequence as in Section 3:

$$a_i = 1 - \frac{2^{i-1} - 1}{2^N - 1}.$$

Then it is easy to see that $1 = a_1 > a_2 > \dots > a_N > 0$ and $2a_{i-1} - a_i > a_1$ for each $2 \leq i \leq N$.

Define functions

$$\begin{aligned} V_1(\mathbf{E}) &= \sum_{i=1}^N V_{1,i} = \sum_{i=1}^N E_i^{a_1 \eta - 1} \\ V_2(\mathbf{E}) &= \sum_{i=1}^{N-1} V_{2,i} = \sum_{i=1}^{N-1} (E_i + E_{i+1})^{a_2 \eta - 1} \\ &\vdots \\ V_n(\mathbf{E}) &= \sum_{k=1}^{N-n+1} V_{n,k} = \sum_{k=1}^{N-n+1} \left(\sum_{j=0}^{n-1} E_{k+j} \right)^{a_n \eta - 1} \end{aligned}$$

$$\begin{aligned} & \vdots \\ V_N(\mathbf{E}) &= V_{N,1} = \left(\sum_{i=1}^N E_i \right)^{a_N \eta^{-1}}, \end{aligned}$$

and

$$V(\mathbf{E}) = \sum_{i=1}^{N-1} V_i(\mathbf{E}).$$

Our aim is to show that $V(\mathbf{E})$ is a Lyapunov function when the value of $V(\mathbf{E})$ is sufficiently large. The excursion time on low energy set then follows from Theorem 3.12.

The main theorem in this section is as follows.

Theorem 4.1. *For any $\eta > 0$ and $h > 0$ small enough, there exist $c_0 > 0$, $M_0 > 1$ depending on η , N , and h , such that*

$$(P^h)V(\mathbf{E}) - V(\mathbf{E}) \leq -c_0 V^\alpha(\mathbf{E})$$

for every $\mathbf{E} \in \{V > M_0\}$, where $\alpha = 1 - \frac{1}{2(1-\eta)}$.

Let $\mathbf{E} = (E_1, \dots, E_N)$ be the initial condition. For the sake of simplicity, we let $R_i = R(E_{i-1}, E_i)$ for all $i = 0 \sim N$. $\mathbf{E}_t$ is given by the following equivalent description: Starting from $t = 0$, a clock rings at an exponentially distributed random time τ_1 with rate $\sum_{i=0}^N R_i$. When the clock rings, one and only one energy exchange occurs between site i and $i + 1$ with probability

$$\frac{R_i}{\sum_{i=0}^N R_i} := R_i / \mathcal{R}.$$

In other words, $\mathbb{P}_{\mathbf{E}}[\mathcal{C}_i(\tau_1)] = R_i / \mathcal{R}$.

Let $\mathbf{E}_{\tau_1^+}$ be the energy configuration immediately after the first energy exchange occurs at τ_1 . We use $V(\mathbf{E}_{\tau_1^+})$ to estimate $P^h V(\mathbf{E})$. The strategy of proving Theorem 4.1 is as follows. In Lemma 4.2 and Corollary 4.3, we show that an energy exchange event can increase a $V_{n,k}$ by at most multiplying a constant. However, if $V_{n,k}$ is sufficiently large and the energy stored at its “boundary site”, i.e., E_{k-1} or E_{k+n} , is sufficiently large, then in Lemma 4.4 we prove that the corresponding energy exchange event will reduce the value of such a $V_{n,k}$ by at least one half. Then in Lemma 4.5 we manage to prove that if $V_{n,k}$ is sufficiently large, the “expected jump” of $V_{n,k}$ at τ_1 can be compensated by the “expected drop” of $V_{n',k'}/4N^2$ for some $V_{n',k'}$. This can always be achieved because $V_{n',k'}$ is significantly greater than $V_{n,k}$ when $V_{n',k'}$ “covers” $V_{n,k}$. Finally, in Lemma 4.6 we proved that the “expected drop” of V at τ_1 is proportional to V^α for $\alpha = 1 - \frac{1}{2(1-\eta)}$. The heuristics of Lemma 4.6 is that for every sufficiently large $V_{n,k}$, the “expected jump” is compensated by $1/4N^2$ of the “expect drop” of some $V_{n',k'}$ (Lemma 4.5). Therefore, at τ_1 the total “expected jump” of those large $V_{n,k}$ s is compensated by $1/4$ of the largest “expected drops” on the left and right side, denoted by V_{n_R,k_R} and V_{n_L,k_L} , respectively. Because the “expected drop” of at least one $V_{n,k}$ is proportional to V^α , we know that the “expected drop” contributed by V_{n_R,k_R} and V_{n_L,k_L} minus the total “expected jump”

of all large $V_{n,k}$ s, is also proportional to V^α . On the other hand, Lemma 4.2 implies the “expected jump” of small $V_{n,k}$ s at τ_1 can be controlled by a constant.

Lemma 4.2. *For any $\mathbf{E} \in \mathbb{R}_+^N$, $\eta > 0$, $1 \leq n \leq N$, and $1 \leq k \leq N - n + 1$,*

$$\mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+})\mathbf{1}_{\mathcal{C}_k(\tau_1)}] \leq \frac{C}{\mathcal{R}} \left(\sum_{i=0}^{n-1} E_{k+i} \right)^{a_n\eta - \frac{1}{2}}$$

and

$$\mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+})\mathbf{1}_{\mathcal{C}_{k+n}(\tau_1)}] \leq \frac{C}{\mathcal{R}} \left(\sum_{i=0}^{n-1} E_{k+i} \right)^{a_n\eta - \frac{1}{2}}$$

for some $C > 0$ depending on N and η .

Proof. At the first energy exchange, if $k \neq 1$ and $k + n \neq N + 1$, we have

$$\begin{aligned} \mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+})\mathbf{1}_{\mathcal{C}_k(\tau_1)}] &= \mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+}) | \mathcal{C}_k(\tau_1)] \mathbb{P}_{\mathbf{E}}[\mathcal{C}_k(\tau_1)] \\ &= \frac{\min\{K, \sqrt{\min\{E_{k-1}, E_k\}}\}}{\mathcal{R}} \int_0^1 [p(E_{k-1} + E_k) + E_{k+1} \cdots + E_{k+n-1}]^{a_n\eta - 1} dp := I_1 \end{aligned}$$

Notice that $a_n\eta$ is small so $a_n\eta - 1 < 0$. If $E_k \geq \frac{1}{2} \sum_{i=1}^{n-1} E_{k+i}$, then

$$I_1 \leq \frac{E_k^{1/2}}{\mathcal{R}} \cdot E_k^{a_n\eta - 1} \int_0^1 p^{a_n\eta - 1} dp \leq \frac{2}{a_n\eta \mathcal{R}} \left(\sum_{i=0}^{n-1} E_{k+i} \right)^{a_n\eta - \frac{1}{2}}.$$

Otherwise,

$$\begin{aligned} I_1 &\leq \frac{E_k^{1/2}}{\mathcal{R}} \cdot (E_{k+1} + \cdots + E_{k+n-1})^{a_n\eta - 1} \\ &\leq 2 \frac{(E_k + \cdots + E_{k+n-1})^{1/2}}{\mathcal{R}} \cdot (E_k + \cdots + E_{k+n-1})^{a_n\eta - 1} \leq \frac{2}{\mathcal{R}} \left(\sum_{i=0}^{n-1} E_{k+i} \right)^{a_n\eta - \frac{1}{2}}. \end{aligned}$$

The same argument holds for

$$\begin{aligned} &\mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+})\mathbf{1}_{\mathcal{C}_{k+n}(\tau_1)}] \\ &= \frac{\min\{K, \sqrt{\min\{E_{k+n-1}, E_{k+n}\}}\}}{\mathcal{R}} \int_0^1 [E_k \cdots + p(E_{k+n-1} + E_{k+n})]^{a_n\eta - 1} dp \end{aligned}$$

by discussing cases $E_{k+n-1} \geq \frac{1}{2} \sum_{i=1}^{n-2} E_{k+i}$ and $E_{k+n-1} < \frac{1}{2} \sum_{i=1}^{n-2} E_{k+i}$.

If $V_{k,n}$ involves the boundary, say $k = 1$, then

$$\begin{aligned} &\mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+})\mathbf{1}_{\mathcal{C}_k(\tau_1)}] \\ &= \frac{\sqrt{\min\{T_L, E_1\}}}{\mathcal{R}} \int_0^\infty \int_0^1 \frac{1}{T_L} [p(x + E_1) + E_2 \cdots + E_n]^{a_n\eta - 1} e^{-x/T_L} dp dx. \end{aligned}$$

It is easy to check that the same argument above still holds. The case of $k+n = N+1$ can be estimated analogously.

The proof is completed by letting $C = \frac{2}{a_n\eta} \geq \frac{2}{a_n\eta}$. $\square$

The calculation in the proof of Lemma 4.2 gives the following corollary.

Corollary 4.3. *For any $\mathbf{E} \in \mathbb{R}_+^N$, $\eta > 0$, $1 \leq n \leq N$, and $1 \leq k \leq N - n + 1$,*

$$\mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+}) \mid \mathcal{C}_k(\tau_1)] \leq CV_{n,k}(\mathbf{E})$$

and

$$\mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+}) \mid \mathcal{C}_{k+n}(\tau_1)] \leq CV_{n,k}(\mathbf{E}),$$

where $C > 0$ is the same as that in Lemma 4.2.

Proof. We have

$$\mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+}) \mid \mathcal{C}_k(\tau_1)] = \int_0^1 [p(E_{k-1} + E_k) + E_{k+1} \cdots + E_{k+n-1}]^{a_n \eta - 1} dp.$$

It follows from the same argument as in the proof of Lemma 4.2 that

$$\begin{aligned} & \int_0^1 [p(E_{k-1} + E_k) + E_{k+1} \cdots + E_{k+n-1}]^{a_n \eta - 1} dp \\ & \leq \max\left\{\frac{2}{a_n \eta}, 2\right\} (E_k + \cdots + E_{k+n-1})^{a_n \eta - 1} := CV_{n,k}(\mathbf{E}). \end{aligned}$$

The proof of

$$\mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+}) \mid \mathcal{C}_{k+n}(\tau_1)] \leq CV_{n,k}(\mathbf{E})$$

is similar. $\square$

Lemma 4.4. *For any $\mathbf{E} \in \mathbb{R}_+^N$ and any $0 < \eta \ll 1$, there exists a constant C' depending on N and η , such that whenever $E_{k+n} > C'(E_k + \cdots + E_{k+n-1})$ (resp. $E_{k-1}(0) > C'(E_k + \cdots + E_{k+n-1})$), we have*

$$\mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+}) \mid \mathcal{C}_{k+n}(\tau_1)] < \frac{1}{2} V_{n,k}(\mathbf{E})$$

(resp.

$$\mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+}) \mid \mathcal{C}_k(\tau_1)] < \frac{1}{2} V_{n,k}(\mathbf{E}).$$

)

Proof. First assume $k + n \neq N + 1$, then we have

$$\begin{aligned} & \mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+}) \mid \mathcal{C}_{k+n}(\tau_1)] \\ & = \int_0^1 (E_k + \cdots + E_{k+n-2} + p(E_{k+n-1} + E_{k+n}))^{a_n \eta - 1} dp \\ & < E_{k+n}^{a_n \eta - 1} \int_0^1 p^{a_n \eta - 1} dp. \end{aligned}$$

If $E_{k+n} > C'(E_k + \cdots + E_{k+n-1})$, we have

$$E_{k+n}^{a_n \eta - 1} \int_0^1 p^{a_n \eta - 1} dp \leq \frac{C'^{a_n \eta - 1}}{a_n \eta} V_{n,k}(\mathbf{E}).$$

To make

$$\frac{C'^{a_n\eta-1}}{a_n\eta} \leq \frac{1}{2},$$

one needs

$$C' \geq \left(\frac{1}{2}a_n\eta\right)^{\frac{1}{a_n\eta-1}}.$$

If $k + n = N + 1$, then

$$\begin{aligned} & \mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+}) | \mathcal{C}_{k+n}(\tau_1)] \\ &= \int_0^1 \int_0^\infty \frac{1}{T_R} (E_k + \dots + E_{k+n-2} + p(E_{k+n-1} + x))^{a_n\eta-1} e^{-x/T_R} dp dx \\ &< \int_0^\infty \frac{1}{T_R} x^{a_n\eta-1} e^{-x/T_R} dx \int_0^1 p^{a_n\eta-1} dp \\ &= T_R^{a_n\eta-1} \frac{\Gamma(a_n\eta)}{a_n\eta}, \end{aligned}$$

where $\Gamma(\cdot)$ is the Gamma function. Therefore to make

$$\mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+}) | \mathcal{C}_{k+n}(\tau_1)] < \frac{1}{2} V_{n,k}(\mathbf{E}),$$

we need

$$V_{n,k}(\mathbf{E}) = (E_k + \dots + E_{k+n-1})^{a_n\eta-1} > 2T_R^{a_n\eta-1} \frac{\Gamma(a_n\eta)}{a_n\eta}.$$

Since $E_{N+1} = T_R$, this is equivalent to

$$E_{N+1} > T_R \left[2T_R^{a_n\eta-1} \cdot \frac{\Gamma(a_n\eta)}{a_n\eta} \right]^{-\frac{1}{a_n\eta-1}} (E_k + \dots + E_{k+n-1}).$$

The case for $\mathcal{C}_k$ is symmetric and can be calculated in the same way. By combining all cases, it is easy to check that if

$$C' = \max_{1 \leq n \leq N} \left\{ \left(\frac{1}{2}a_n\eta\right)^{\frac{1}{a_n\eta-1}}, T_R \left[2T_R^{a_n\eta-1} \cdot \frac{\Gamma(a_n\eta)}{a_n\eta} \right]^{-\frac{1}{a_n\eta-1}}, T_L \left[2T_L^{a_n\eta-1} \cdot \frac{\Gamma(a_n\eta)}{a_n\eta} \right]^{-\frac{1}{a_n\eta-1}} \right\},$$

we have the desired property for all n and k .

□

Lemma 4.5. *For any $\mathbf{E} \in \mathbb{R}_+^N$ and any $0 < \eta \ll 1$, there exists an $M < \infty$ depending on η and N , such that for any $V_{n,k}(\mathbf{E}) > M$, if*

$$\mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+}) \mathbf{1}_{\mathcal{C}_{k+n}(\tau_1)}] \geq V_{n,k}(\mathbf{E}) \mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k+n}(\tau_1)]$$

(resp.

$$\mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+}) \mathbf{1}_{\mathcal{C}_k(\tau_1)}] \geq V_{n,k}(\mathbf{E}) \mathbb{P}_{\mathbf{E}}[\mathcal{C}_k(\tau_1)],$$

) then there exists k' and n' , such that

$$\begin{aligned} & \mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+}) \mathbf{1}_{\mathcal{C}_{k+n}(\tau_1)}] - V_{n,k}(\mathbf{E}) \mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k+n}(\tau_1)] \\ & \leq \frac{1}{4N^2} \left\{ V_{n',k'}(\mathbf{E}) \mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k'+n'}(\tau_1)] - \mathbb{E}_{\mathbf{E}}[V_{n',k'}(\mathbf{E}_{\tau_1^+}) \mathbf{1}_{\mathcal{C}_{k'+n'}(\tau_1)}] \right\}. \end{aligned}$$

(resp.

$$\begin{aligned} & \mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+})\mathbf{1}_{\mathcal{C}_k(\tau_1)}] - V_{n,k}(\mathbf{E})\mathbb{P}_{\mathbf{E}}[\mathcal{C}_k(\tau_1)] \\ & \leq \frac{1}{4N^2} \left\{ V_{n',k'}(\mathbf{E})\mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k'}(\tau_1)] - \mathbb{E}_{\mathbf{E}}[V_{n',k'}(\mathbf{E}_{\tau_1^+})\mathbf{1}_{\mathcal{C}_{k'}(\tau_1)}] \right\}. \end{aligned}$$

)

Proof. By symmetry, it is sufficient to consider the case of

$$\mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+})\mathbf{1}_{\mathcal{C}_{k+n}(\tau_1)}] \geq V_{n,k}(\mathbf{E})\mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k+n}(\tau_1)].$$

Let $k' = k$ and n' be the first $n' > n - 1$ such that

$$E_{k+n'} > C'(E_k + \dots + E_{k+n'-1})$$

where C' is the constant defined in Lemma 4.4. When M is large, the sum $E_k + \dots + E_{k+n'-1}$ is small. Note that $E_{N+1} = T_L$. Hence when M is sufficiently large, for any $V_{n,k}(\mathbf{E}_0) > M$, one can always find such an n' .

By Lemma 4.4, if $n' = n$, the energy exchange event $\mathcal{C}_{k+n}(\tau_1)$ will only bring the expected value of $V_{n,k}$ down. Therefore, it is sufficient to consider the case of $n' \geq n + 1$. The lemma follows if one can prove either

$$\begin{aligned} & \mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+})\mathbf{1}_{\mathcal{C}_{k+n}(\tau_1)}] - V_{n,k}(\mathbf{E})\mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k+n}(\tau_1)] \\ & \leq \frac{1}{4N^2} \left\{ V_{n',k}(\mathbf{E})\mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k+n'}(\tau_1)] - \mathbb{E}_{\mathbf{E}}[V_{n',k}(\mathbf{E}_{\tau_1^+})\mathbf{1}_{\mathcal{C}_{k+n'}(\tau_1)}] \right\} := \frac{1}{4N^2} I_1 \end{aligned}$$

or

$$\begin{aligned} & \mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+})\mathbf{1}_{\mathcal{C}_{k+n}(\tau_1)}] - V_{n,k}(\mathbf{E})\mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k+n}(\tau_1)] \\ & \leq \frac{1}{4N^2} \left\{ V_{1,k+n'-1}(\mathbf{E})\mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k+n'}(\tau_1)] - \mathbb{E}_{\mathbf{E}}[V_{1,k+n'-1}(\mathbf{E}_{\tau_1^+})\mathbf{1}_{\mathcal{C}_{k+n'}(\tau_1)}] \right\} := \frac{1}{4N^2} I_2 \end{aligned}$$

Let $x = E_k + \dots + E_{k+n'-2}$, $y = E_{k+n'-1}$, and $z = E_{k+n'}$. Since $a_{n'} < a_n$, if x is sufficiently small, by Lemma 4.2,

$$\begin{aligned} \mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+})\mathbf{1}_{\mathcal{C}_{k+n}(\tau_1)}] & \leq \frac{C}{\mathcal{R}}(E_k + \dots + E_{k+n-1})^{a_n\eta - \frac{1}{2}} \\ & \leq \frac{C}{\mathcal{R}} \left(\frac{1}{NC'^{N-1}}(E_k + \dots + E_{k+n'-2}) \right)^{a_n\eta - \frac{1}{2}} \\ & \leq \frac{1}{\mathcal{R}} x^{a_{n'-1}\eta - \frac{1}{2}} \cdot \left(C \cdot C'^{(\frac{1}{2} - a_n\eta)(N-1)} \cdot N^{\frac{1}{2} - a_n\eta} \cdot x^{(a_n - a_{n'-1})\eta} \right) \\ & \leq \frac{1}{\mathcal{R}} x^{a_{n'-1}\eta - \frac{1}{2}}, \end{aligned}$$

where C is the constant in Lemma 4.2. The second inequality above follows from

$$E_{k+m} \leq C'(E_k + \dots + E_{k+m-1})$$

for each $n \leq m < n'$.

The requirement of small x can be satisfied by making $M = M(\eta)$ sufficiently large, because

$$x \leq NC'^{N-1}(E_k + \dots + E_{k+n-1}) \leq NC'^{N-1}M^{\frac{1}{a_n\eta - 1}}.$$

On the other hand, notice that K is sufficiently large and $y < z$ by the assumption of n' (Obviously C' in Lemma 4.4 is greater than 1). Therefore $R(E_{k+n'-1}, E_{k+n'}) = \sqrt{y}$. Hence we have

$$I_1 = \left\{ (x+y)^{a_{n'}\eta-1} - \int_0^1 [x+p(y+z)]^{a_{n'}\eta-1} dp \right\} \frac{\sqrt{y}}{\mathcal{R}}$$

and

$$I_2 = \left\{ y^{a_1\eta-1} - \int_0^1 [p(y+z)]^{a_1\eta-1} dp \right\} \frac{\sqrt{y}}{\mathcal{R}}.$$

Since $z > C'(y+x) > C'y$ and $y < C'x$, by Lemma 4.4, we have

$$I_1 \geq \frac{1}{2}(x+y)^{a_{n'}\eta-1} \frac{\sqrt{y}}{\mathcal{R}} \geq \frac{1}{2(1+C')} \frac{x^{a_{n'}\eta-1} \sqrt{y}}{\mathcal{R}}$$

and

$$I_2 \geq \frac{1}{2} y^{a_1\eta-1} \frac{\sqrt{y}}{\mathcal{R}}.$$

We claim that

$$\max \left\{ \frac{1}{2} y^{a_1\eta-\frac{1}{2}}, \frac{1}{2C'} x^{a_{n'}\eta-1} \sqrt{y} \right\} > 4N^2 x^{a_{n'-1}\eta-\frac{1}{2}}.$$

It is easy to see that the lemma follows from this claim.

Proof of the claim: Let $0 < \epsilon \ll 1$ be a small constant depending on $\{a_n\}_{n=1}^N$.

Case 1: $y < x^{1+2(a_{n'-1}-a_{n'})\eta-\epsilon}$, then

$$\frac{1}{2} y^{a_1\eta-\frac{1}{2}} > \frac{1}{2} x^{(a_1\eta-\frac{1}{2})[1+2(a_{n'-1}-a_{n'})\eta-\epsilon]}.$$

Note that $2a_{n'-1} - a_{n'} > a_1$ and $\eta \ll 1$, we have

$$\begin{aligned} & (a_1\eta - \frac{1}{2})[1 + 2(a_{n'-1} - a_{n'})\eta - \epsilon] \\ = & -\frac{1}{2} + a_{n'-1}\eta - \epsilon + (a_1 + a_{n'} - 2a_{n'-1})\eta + \epsilon + \epsilon(\frac{1}{2} - a_1\eta) + 2a_1(a_{n'-1} - a_{n'})\eta^2 \\ < & -\frac{1}{2} + a_{n'-1}\eta - \epsilon \end{aligned}$$

if

$$(2a_{n'-1} - a_{n'} - a_1)\eta > \frac{3}{2}\epsilon + 2a_1(a_{n'-1} - a_{n'})\eta^2,$$

which can be achieved by making ϵ and η sufficiently small.

Therefore,

$$\frac{1}{2} y^{a_1\eta-\frac{1}{2}} > \frac{1}{2} x^{-\epsilon} \cdot x^{a_{n'}\eta-1} > 4N^2 x^{a_{n'-1}\eta-\frac{1}{2}}$$

when x is sufficiently small, which can be made by letting M large enough.

Case 2: $y \geq x^{1+2(a_{n'-1}-a_{n'})\eta-\epsilon}$, then let ϵ be as in **Case 1**, we have

$$\begin{aligned} \frac{1}{2C'} x^{a_{n'}\eta-1} \sqrt{y} &\geq \frac{1}{2C'} x^{a_{n'}\eta-1} \cdot x^{\frac{1}{2}+a_{n'-1}\eta-a_{n'}\eta-\frac{\epsilon}{2}} \\ &= \frac{1}{2C'} x^{a_{n'-1}\eta-\frac{1}{2}} \cdot x^{-\frac{\epsilon}{2}} \\ &> 4N^2 x^{a_{n'-1}\eta-\frac{1}{2}} \end{aligned}$$

if x is sufficiently small. Again, this can be achieved by letting M sufficiently large. $\square$

Lemma 4.6. *For any $\mathbf{E} \in \mathbb{R}_+^N$ and any $0 < \eta \ll 1$, there exist constants $\alpha = 1 - \frac{1}{2(1-\eta)}$, C_1 and M' depending on N and η , such that*

$$\mathbb{E}_{\mathbf{E}}[V(\mathbf{E}_{\tau_1^+})] \leq V(\mathbf{E}) - \frac{C_1}{\mathcal{R}} V^\alpha(\mathbf{E})$$

for all $V(\mathbf{E}) > M'$.

Proof. Step 1. First we show that there exists $\tilde{n}, \tilde{k}$ such that the “expected drop” of $V_{\tilde{n}, \tilde{k}}$ at τ_1 is proportional to $V^\alpha(\mathbf{E})$. Let $V_{\tilde{n}, \tilde{k}}(\mathbf{E})$ be the maximum of $\{V_{i,j}(\mathbf{E})\}$. Therefore $V_{\tilde{n}, \tilde{k}}(\mathbf{E}) \geq \frac{1}{N(N+1)} V(\mathbf{E})$. Since $V_{\tilde{n}, \tilde{k}}(\mathbf{E})$ is the maximum, we have

$$(E_{\tilde{k}} + \cdots + E_{\tilde{k}+\tilde{n}-1})^{a_{\tilde{n}}\eta-1} > (E_{\tilde{k}-1} + \cdots + E_{\tilde{k}+\tilde{n}-1})^{a_{\tilde{n}+1}\eta-1}$$

and

$$(E_{\tilde{k}} + \cdots + E_{\tilde{k}+\tilde{n}-1})^{a_{\tilde{n}}\eta-1} > (E_{\tilde{k}} + \cdots + E_{\tilde{k}+\tilde{n}})^{a_{\tilde{n}+1}\eta-1}.$$

Since $a_{\tilde{n}+1} < a_{\tilde{n}}$, when M' is sufficiently large, we have

$$E_{\tilde{k}-1} > C'(E_{\tilde{k}} + \cdots + E_{\tilde{k}+\tilde{n}-1})$$

and

$$E_{\tilde{k}+\tilde{n}} > C'(E_{\tilde{k}} + \cdots + E_{\tilde{k}+\tilde{n}-1}),$$

where C' is as in Lemma 4.4. In addition, we have

$$E_{\tilde{k}}^{a_1\eta-1} \leq (E_{\tilde{k}} + \cdots + E_{\tilde{k}+\tilde{n}-1})^{a_{\tilde{n}}\eta-1}$$

and

$$E_{\tilde{k}+\tilde{n}-1}^{a_1\eta-1} \leq (E_{\tilde{k}} + \cdots + E_{\tilde{k}+\tilde{n}-1})^{a_{\tilde{n}}\eta-1}.$$

Since M' is assumed to be sufficiently large, we have $\min\{K, \min\{E_{\tilde{k}-1}, E_{\tilde{k}}\}\} = E_{\tilde{k}}$ and $\min\{K, \min\{E_{\tilde{k}+\tilde{n}-1}, E_{\tilde{k}+\tilde{n}}\}\} = E_{\tilde{k}+\tilde{n}-1}$. Therefore,

$$\begin{aligned} &\mathbb{E}_{\mathbf{E}}[V_{\tilde{n}, \tilde{k}}(\mathbf{E}_{\tau_1^+})] - V_{\tilde{n}, \tilde{k}}(\mathbf{E}) \\ &= \frac{\sqrt{E_{\tilde{k}}}}{\mathcal{R}} \left\{ \int_0^1 [p(E_{\tilde{k}-1} + E_{\tilde{k}}) + \cdots + E_{\tilde{n}+\tilde{k}-1}]^{a_{\tilde{n}}\eta-1} dp - (E_{\tilde{k}} + \cdots + E_{\tilde{k}+\tilde{n}-1})^{a_{\tilde{n}}\eta-1} \right\} \\ &+ \frac{\sqrt{E_{\tilde{k}+\tilde{n}-1}}}{\mathcal{R}} \left\{ \int_0^1 [E_{\tilde{k}} + \cdots + p(E_{\tilde{n}+\tilde{k}-1} + E_{\tilde{n}+\tilde{k}})]^{a_{\tilde{n}}\eta-1} dp - (E_{\tilde{k}} + \cdots + E_{\tilde{k}+\tilde{n}-1})^{a_{\tilde{n}}\eta-1} \right\} \\ &:= I_1 + I_2. \end{aligned}$$

It follows from the same calculation as in Lemma 4.4 that

$$(4.1) \quad \begin{aligned} I_i &\leq -\frac{1}{2}(E_{\tilde{k}} + \cdots + E_{\tilde{n}+\tilde{k}-1})^{a_{\tilde{n}}\eta-1} \cdot (E_{\tilde{k}} + \cdots + E_{\tilde{n}+\tilde{k}-1})^{\frac{a_{\tilde{n}}\eta-1}{2(a_1\eta-1)}} \cdot \frac{1}{\mathcal{R}} \\ &= -\frac{1}{2\mathcal{R}}V_{\tilde{n},\tilde{k}}^\alpha(\mathbf{E}) \end{aligned}$$

for $i = 1, 2$, where

$$\alpha = 1 - \frac{1}{2(1 - a_1\eta)}.$$

Step 2. Let M be as in Lemma 4.5. Let $V_{n_1,k_1}, \dots, V_{n_m,k_m}$ be functions that satisfy $V_{n_i,k_i}(\mathbf{E}) \geq M$. The aim of this step is to prove that the total “expected jump” of these V_{n_i,k_i} at τ_1 is dominated by the “expected drop” of *some* V_{n_R,k_R} and V_{n_L,k_L} from left and right side, respectively.

By Lemma 4.5, we can construct sequences $\{(\hat{n}'_i, \hat{k}'_i)\}_{i=1}^m$ and $\{(\hat{n}''_i, \hat{k}''_i)\}_{i=1}^m$ such that if

$$\mathbb{E}_{\mathbf{E}}[V_{n_i,k_i}(\mathbf{E}_{\tau_1^+}), \mathbf{1}_{\mathcal{C}_{k_i+n_i}(\tau_1)}] \geq V_{n_i,k_i}(\mathbf{E})\mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k_i+n_i}(\tau_1)],$$

then

$$\begin{aligned} &\mathbb{E}_{\mathbf{E}}[V_{n_i,k_i}(\mathbf{E}_{\tau_1^+}), \mathbf{1}_{\mathcal{C}_{k_i+n_i}(\tau_1)}] - V_{n_i,k_i}(\mathbf{E})\mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k_i+n_i}(\tau_1)] \\ &\leq \frac{1}{4N^2} \{V_{\hat{n}'_i, \hat{k}'_i}(\mathbf{E})\mathbb{P}_{\mathbf{E}}[\mathcal{C}_{\hat{k}'_i+\hat{n}'_i}(\tau_1)] - \mathbb{E}_{\mathbf{E}}[V_{\hat{n}'_i, \hat{k}'_i}(\mathbf{E}_{\tau_1^+}), \mathbf{1}_{\mathcal{C}_{\hat{k}'_i+\hat{n}'_i}(\tau_1)}]\} \end{aligned}$$

and if

$$\mathbb{E}_{\mathbf{E}}[V_{n_i,k_i}(\mathbf{E}_{\tau_1^+}), \mathbf{1}_{\mathcal{C}_{k_i}(\tau_1)}] \geq V_{k_i,n_i}(\mathbf{E})\mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k_i}(\tau_1)],$$

then

$$\begin{aligned} &\mathbb{E}_{\mathbf{E}}[V_{n_i,k_i}(\mathbf{E}_{\tau_1^+}), \mathbf{1}_{\mathcal{C}_{k_i}(\tau_1)}] - V_{k_i,n_i}(\mathbf{E})\mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k_i}(\tau_1)] \\ &\leq \frac{1}{4N^2} \{V_{\hat{n}''_i, \hat{k}''_i}(\mathbf{E})\mathbb{P}_{\mathbf{E}}[\mathcal{C}_{\hat{k}''_i}(\tau_1)] - \mathbb{E}_{\mathbf{E}}[V_{\hat{n}''_i, \hat{k}''_i}(\mathbf{E}_{\tau_1^+}), \mathbf{1}_{\mathcal{C}_{\hat{k}''_i}(\tau_1)}]\}. \end{aligned}$$

When conditions

$$\mathbb{E}_{\mathbf{E}}[V_{n_i,k_i}(\mathbf{E}_{\tau_1^+}), \mathbf{1}_{\mathcal{C}_{k_i+n_i}(\tau_1)}] \geq V_{n_i,k_i}(\mathbf{E})\mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k_i+n_i}(\tau_1)],$$

(resp.

$$\mathbb{E}_{\mathbf{E}}[V_{n_i,k_i}(\mathbf{E}_{\tau_1^+}), \mathbf{1}_{\mathcal{C}_{k_i}(\tau_1)}] \geq V_{k_i,n_i}(\mathbf{E})\mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k_i}(\tau_1)],$$

) is not satisfied, we simply let $\hat{n}'_i = n_i, \hat{k}'_i = k_i$ (resp. $\hat{n}''_i = n_i, \hat{k}''_i = k_i$). Therefore, the “expected jump” of V_{n_i,k_i} at τ_1 is dominated by the right and left “expected drop” of $V_{\hat{n}'_i, \hat{k}'_i}$ and $V_{\hat{n}''_i, \hat{k}''_i}$, respectively.

Let $V_{n_R,k_R}(\mathbf{E})$ be the function with the biggest “right drop” at τ_1 , i.e.,

$$\begin{aligned} &V_{n,k}(\mathbf{E})\mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k+n}(\tau_1)] - \mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+}), \mathbf{1}_{\mathcal{C}_{k+n}(\tau_1)}] \\ &\leq V_{n_R,k_R}(\mathbf{E})\mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k_R+n_R}(\tau_1)] - \mathbb{E}_{\mathbf{E}}[V_{n_R,k_R}(\mathbf{E}_{\tau_1^+}), \mathbf{1}_{\mathcal{C}_{k_R+n_R}(\tau_1)}] \end{aligned}$$

for all n, k , and $V_{n_L,k_L}(\mathbf{E})$ be the function with the biggest “left drop”, i.e.

$$\begin{aligned} &V_{n,k}(\mathbf{E})\mathbb{P}_{\mathbf{E}}[\mathcal{C}_k(\tau_1)] - \mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+}), \mathbf{1}_{\mathcal{C}_k(\tau_1)}] \\ &\leq V_{n_L,k_L}(\mathbf{E})\mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k_L}(\tau_1)] - \mathbb{E}_{\mathbf{E}}[V_{n_L,k_L}(\mathbf{E}_{\tau_1^+}), \mathbf{1}_{\mathcal{C}_{k_L}(\tau_1)}] \end{aligned}$$

for all n, k . Both terms should be negative when M' is sufficiently large because of the argument about $V_{\tilde{n}, \tilde{k}}$ in **Step 1**.

Since there is only $\frac{1}{2}N(N+1)$ $V_{n,k}$ s, we have $m < \frac{1}{2}N(N+1)$. Hence

$$\begin{aligned}
& \sum_{i=1}^m \mathbb{E}_{\mathbf{E}}[V_{n_i, k_i}(\mathbf{E}_{\tau_1^+})] - V_{n_i, k_i}(\mathbf{E}) \\
&= \sum_{i=1}^m \{ \mathbb{E}_{\mathbf{E}}[V_{n_i, k_i}(\mathbf{E}_{\tau_1^+}) \mathbf{1}_{\mathcal{C}_{k_i+n_i}(\tau_1)}] - V_{n_i, k_i}(\mathbf{E}) \mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k_i+n_i}(\tau_1)] \\
&\quad + \mathbb{E}_{\mathbf{E}}[V_{n_i, k_i}(\mathbf{E}_{\tau_1^+}) \mathbf{1}_{\mathcal{C}_{k_i}(\tau_1)}] - V_{n_i, k_i}(\mathbf{E}) \mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k_i}(\tau_1)] \} \\
&\leq \frac{1}{4N^2} \sum_{i=1}^m \left\{ \min \{ \mathbb{E}_{\mathbf{E}}[V_{\hat{n}'_i, \hat{k}'_i}(\mathbf{E}_{\tau_1^+}) \mathbf{1}_{\mathcal{C}_{\hat{k}'_i+\hat{n}'_i}(\tau_1)}] - V_{\hat{n}'_i, \hat{k}'_i}(\mathbf{E}) \mathbb{P}_{\mathbf{E}}[\mathcal{C}_{\hat{k}'_i+\hat{n}'_i}(\tau_1)], 0 \} \right. \\
&\quad \left. + \min \{ \mathbb{E}_{\mathbf{E}}[V_{\hat{n}''_i, \hat{k}''_i}(\mathbf{E}_{\tau_1^+}) \mathbf{1}_{\mathcal{C}_{\hat{k}''_i}(\tau_1)}] - V_{\hat{n}''_i, \hat{k}''_i}(\mathbf{E}) \mathbb{P}_{\mathbf{E}}[\mathcal{C}_{\hat{k}''_i}(\tau_1)], 0 \} \right\} \\
&\leq \frac{1}{4} \{ V_{n_R, k_R}(\mathbf{E}) \mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k_R+n_R}(\tau_1)] - \mathbb{E}_{\mathbf{E}}[V_{n_R, k_R}(\mathbf{E}_{\tau_1^+}) \mathbf{1}_{\mathcal{C}_{k_R+n_R}(\tau_1)}] \\
&\quad + V_{n_L, k_L}(\mathbf{E}) \mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k_L}(\tau_1)] - \mathbb{E}_{\mathbf{E}}[V_{n_L, k_L}(\mathbf{E}_{\tau_1^+}) \mathbf{1}_{\mathcal{C}_{k_L}(\tau_1)}] \}.
\end{aligned}$$

Step 3. Then we can finalize the entire proof. We have

$$\begin{aligned}
& \mathbb{E}_{\mathbf{E}}(V(\mathbf{E}_{\tau_1^+})) - V(\mathbf{E}) \\
&= \sum_{n,k} \{ \mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+}) \mathbf{1}_{\mathcal{C}_{k+n}(\tau_1)}] - V_{n,k}(\mathbf{E}) \mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k+n}(\tau_1)] \\
&\quad + \mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+}) \mathbf{1}_{\mathcal{C}_k(\tau_1)}] - V_{n,k}(\mathbf{E}) \mathbb{P}_{\mathbf{E}}[\mathcal{C}_k(\tau_1)] \} \\
&= \sum_{i=1}^m \{ \mathbb{E}_{\mathbf{E}}[V_{n_i, k_i}(\mathbf{E}_{\tau_1^+}) \mathbf{1}_{\mathcal{C}_{k_i+n_i}(\tau_1)}] - V_{n_i, k_i}(\mathbf{E}) \mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k_i+n_i}(\tau_1)] \\
&\quad + \mathbb{E}_{\mathbf{E}}[V_{n_i, k_i}(\mathbf{E}_{\tau_1^+}) \mathbf{1}_{\mathcal{C}_{k_i}(\tau_1)}] - V_{n_i, k_i}(\mathbf{E}) \mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k_i}(\tau_1)] \} \\
&\quad + \sum_{k,n \text{ } V_{n,k} < M} \{ \mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+})] - V_{n,k}(\mathbf{E}) \} \\
&\leq \frac{3}{4} \{ \mathbb{E}_{\mathbf{E}}[V_{n_R, k_R}(\mathbf{E}_{\tau_1^+}) \mathbf{1}_{\mathcal{C}_{n_R+k_R}(\tau_1)}] - V_{n_R, k_R}(\mathbf{E}) \mathbb{P}_{\mathbf{E}}[\mathcal{C}_{n_R+k_R}(\tau_1)] \\
&\quad + \mathbb{E}_{\mathbf{E}}[V_{n_L, k_L}(\mathbf{E}_{\tau_1^+}) \mathbf{1}_{\mathcal{C}_{k_L}(\tau_1)}] - V_{n_L, k_L}(\mathbf{E}) \mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k_L}(\tau_1)] \} \\
&\quad + \sum_{k,n \text{ } V_{n,k} < M} \{ \mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+})] - V_{n,k}(\mathbf{E}) \} \\
&\leq \frac{3}{4}(I_3 + I_4) + I_5,
\end{aligned}$$

where

$$\begin{aligned}
I_3 &= \mathbb{E}_{\mathbf{E}}[V_{n_R, k_R}(\mathbf{E}_{\tau_1^+}) \mathbf{1}_{\mathcal{C}_{n_R+k_R}(\tau_1)}] - V_{n_R, k_R}(\mathbf{E}) \mathbb{P}_{\mathbf{E}}[\mathcal{C}_{n_R+k_R}(\tau_1)], \\
I_4 &= \mathbb{E}_{\mathbf{E}}[V_{n_L, k_L}(\mathbf{E}_{\tau_1^+}) \mathbf{1}_{\mathcal{C}_{k_L}(\tau_1)}] - V_{n_L, k_L}(\mathbf{E}) \mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k_L}(\tau_1)],
\end{aligned}$$

and

$$I_5 = \sum_{k,n} \sum_{V_{n,k} < M} \{ \mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+})] - V_{n,k}(\mathbf{E}) \}.$$

It follows from Corollary 4.3 that

$$\begin{aligned} I_5 &= \sum_{k,n} \sum_{V_{n,k} < M} \{ \mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+}) \mathbf{1}_{\mathcal{C}_{k+n}(\tau_1)}] - V_{n,k}(\mathbf{E}) \mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k+n}(\tau_1)] \\ &\quad + \mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+}), \mathbf{1}_{\mathcal{C}_k(\tau_1)}] - V_{n,k}(\mathbf{E}) \mathbb{P}_{\mathbf{E}}[\mathcal{C}_k(\tau_1)] \} \\ &\leq \sum_{k,n} \sum_{V_{n,k} < M} \{ \mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+}) | \mathcal{C}_{k+n}(\tau_1)] \mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k+n}(\tau_1)] + \mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+}) | \mathcal{C}_k(\tau_1)] \mathbb{P}_{\mathbf{E}}[\mathcal{C}_k(\tau_1)] \} \\ &\leq \sum_{k,n} \sum_{V_{n,k} < M} \{ \mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+}) | \mathcal{C}_{k+n}(\tau_1)] + \mathbb{E}_{\mathbf{E}}[V_{n,k}(\mathbf{E}_{\tau_1^+}) | \mathcal{C}_k(\tau_1)] \} \cdot \frac{2K}{\mathcal{R}} \\ &\leq N(N+1)CMK \cdot \frac{1}{\mathcal{R}}. \end{aligned}$$

In addition, by the definition of n_R, k_R and n_L, k_L , we have

$$\begin{aligned} &V_{\tilde{n}, \tilde{k}}(\mathbf{E}) \mathbb{P}_{\mathbf{E}}[\mathcal{C}_{\tilde{k}+\tilde{n}}(\tau_1)] - \mathbb{E}_{\mathbf{E}}[V_{\tilde{n}, \tilde{k}}(\mathbf{E}_{\tau_1^+}) \mathbf{1}_{\mathcal{C}_{\tilde{k}+\tilde{n}}(\tau_1)}] \\ &\leq V_{n_R, k_R}(\mathbf{E}) \mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k_R+n_R}(\tau_1)] - \mathbb{E}_{\mathbf{E}}[V_{n_R, k_R}(\mathbf{E}_{\tau_1^+}) \mathbf{1}_{\mathcal{C}_{k_R+n_R}(\tau_1)}] \end{aligned}$$

and

$$\begin{aligned} &V_{\tilde{n}, \tilde{k}}(\mathbf{E}) \mathbb{P}_{\mathbf{E}}[\mathcal{C}_{\tilde{k}}(\tau_1)] - \mathbb{E}_{\mathbf{E}}[V_{\tilde{n}, \tilde{k}}(\mathbf{E}_{\tau_1^+}) \mathbf{1}_{\mathcal{C}_{\tilde{k}}(\tau_1)}] \\ &\leq V_{n_L, k_L}(\mathbf{E}) \mathbb{P}_{\mathbf{E}}[\mathcal{C}_{k_L}(\tau_1)] - \mathbb{E}_{\mathbf{E}}[V_{n_L, k_L}(\mathbf{E}_{\tau_1^+}) \mathbf{1}_{\mathcal{C}_{k_L}(\tau_1)}], \end{aligned}$$

where $\tilde{n}$ and $\tilde{k}$ are from **Step 1**.

By inequality (4.1) in **Step 1**, we have

$$\begin{aligned} I_3 + I_4 &\leq (\mathbb{E}_{\mathbf{E}}[V_{\tilde{n}, \tilde{k}}(\mathbf{E}_{\tau_1^+}) \mathbf{1}_{\mathcal{C}_{\tilde{k}+\tilde{n}}(\tau_1)}] - V_{\tilde{n}, \tilde{k}}(\mathbf{E}) \mathbb{P}_{\mathbf{E}}[\mathcal{C}_{\tilde{k}+\tilde{n}}(\tau_1)]) \\ &\quad + (\mathbb{E}_{\mathbf{E}}[V_{\tilde{n}, \tilde{k}}(\mathbf{E}_{\tau_1^+}) \mathbf{1}_{\mathcal{C}_{\tilde{k}}(\tau_1)}] - V_{\tilde{n}, \tilde{k}}(\mathbf{E}) \mathbb{P}_{\mathbf{E}}[\mathcal{C}_{\tilde{k}}(\tau_1)]) \\ &= \mathbb{E}_{\mathbf{E}}[V_{\tilde{n}, \tilde{k}}(\mathbf{E}_{\tau_1^+})] - V_{\tilde{n}, \tilde{k}}(\mathbf{E}) \\ &\leq -\frac{1}{\mathcal{R}} V_{\tilde{n}, \tilde{k}}^{\alpha}(\mathbf{E}) \leq -\frac{1}{\mathcal{R}} \cdot \left(\frac{1}{N(N+1)} \right)^{\alpha} V^{\alpha}(\mathbf{E}). \end{aligned}$$

One can then further choose $M' > M$ such that

$$\frac{1}{4\mathcal{R}} \cdot \left(\frac{1}{N(N+1)} \right)^{\alpha} M'^{\alpha} > I_5.$$

Therefore, for every $\mathbf{E}$ such that $V(\mathbf{E}) > M'$, we have

$$\mathbb{E}_{\mathbf{E}}(V(\mathbf{E}_{\tau_1^+}) | \mathbf{E}) - V(\mathbf{E}) \leq -\frac{1}{\mathcal{R}} \cdot \frac{1}{2} \left(\frac{1}{N(N+1)} \right)^{\alpha} V^{\alpha}(\mathbf{E}) := -\frac{C_1}{\mathcal{R}} V^{\alpha}(\mathbf{E}).$$

This completes the proof. □

Proof of Theorem 4.1.

Let $0 = \tau_0 = \tau_1 < \tau_2 < \dots$ be the times of clock rings. Let $B_0 = \{\mathbf{E} \mid V(\mathbf{E}) > M'\}$ where M' is the constant in Lemma 4.6. By the Markov property and Lemma 4.6, for any $\mathbf{E} \in \mathbb{R}_+^N$ we have

$$(4.2) \quad \mathbb{E}_{\mathbf{E}}[V(\mathbf{E}_{\tau_{n+1}^+}) \mid \mathbf{E}_{\tau_n^+}] = \mathbb{E}_{\mathbf{E}_{\tau_n^+}}[V(\mathbf{E}_{\tau_1^+})] \leq V(\mathbf{E}_{\tau_n^+}) - \frac{C_1}{\mathcal{R}} V(\mathbf{E}_{\tau_n^+})^\alpha$$

if $\mathbf{E}_{\tau_n^+} \in B_0$. Otherwise, for each $\mathbf{E}_{\tau_n^+} \notin B_0$, by the Markov property we have the uniform bound

$$(4.3) \quad \begin{aligned} & \mathbb{E}_{\mathbf{E}}[V(\mathbf{E}_{\tau_{n+1}^+}) \mid \mathbf{E}_{\tau_n^+}] \\ &= \mathbb{E}_{\mathbf{E}_{\tau_n^+}}[V_{n,k}(\mathbf{E}_{\tau_1^+}) \mid \mathcal{C}_k(\tau_1)] \mathbb{P}_{\mathbf{E}_{\tau_n^+}}[\mathcal{C}_k(\tau_1)] + \mathbb{E}_{\mathbf{E}_{\tau_n^+}}[V_{n,k}(\mathbf{E}_{\tau_1^+}) \mid \mathcal{C}_{k+n}(\tau_1)] \mathbb{P}_{\mathbf{E}_{\tau_n^+}}[\mathcal{C}_{k+n}(\tau_1)] \\ &\leq \mathbb{E}_{\mathbf{E}_{\tau_n^+}}[V_{n,k}(\mathbf{E}_{\tau_1^+}) \mid \mathcal{C}_k(\tau_1)] + \mathbb{E}_{\mathbf{E}_{\tau_n^+}}[V_{n,k}(\mathbf{E}_{\tau_1^+}) \mid \mathcal{C}_{k+n}(\tau_1)] \\ &\leq 2CV_{n,k}(\mathbf{E}_{\tau_n^+}) \leq 2CM', \end{aligned}$$

where C is the constant in Lemma 4.2.

Let $S = \inf\{n, \tau_n > h\}$, and define $\hat{\tau}_n = \min\{\tau_n, \tau_{S-1}\}$. Then

$$P^h V(\mathbf{E}) = \lim_{n \rightarrow \infty} \mathbb{E}_{\mathbf{E}}[V(\mathbf{E}_{\hat{\tau}_n^+}) \mathbf{1}_{\{S \leq n+1\}}] \leq \lim_{n \rightarrow \infty} \mathbb{E}_{\mathbf{E}}[V(\mathbf{E}_{\hat{\tau}_n^+})].$$

We will prove a uniform bound for $\mathbb{E}_{\mathbf{E}}[V(\mathbf{E}_{\hat{\tau}_n^+})]$. By assuming the worse of (4.2) and (4.3), we have

$$\mathbb{E}_{\mathbf{E}}[V(\mathbf{E}_{\hat{\tau}_{n+1}^+}) \mid \tau_{n+1} \leq h] \leq \mathbb{E}_{\mathbf{E}}[V(\mathbf{E}_{\hat{\tau}_n^+}) \mid \tau_{n+1} \leq h] + CM'$$

for every $n \geq 0$. Notice that for given $\mathbf{E}_{\tau_n^+}$, $\mathbf{E}_{\tau_{n+1}^+}$ is independent of $\tau_{n+1} - \tau_n$. Therefore conditioning on $\tau_{n+1} \leq h$ does not affect the bounds in (4.2) and (4.3). Since $\mathcal{R} \leq NK$, for each $\mathbf{E}_{\tau_n^+}$, we have

$$\mathbb{P}[\tau_{n+1} \leq h \mid \mathbf{E}_{\tau_n^+}, \tau_n \leq h] \leq (1 - e^{-hNK}).$$

Therefore, inductively we have

$$\mathbb{P}_{\mathbf{E}}[\tau_{n+1} \leq h] \leq (1 - e^{-hNK})^{n+1}$$

for all $n \geq 0$. This implies

$$(4.4) \quad \begin{aligned} & \mathbb{E}_{\mathbf{E}}[V(\mathbf{E}_{\hat{\tau}_{n+1}^+})] \\ &= \mathbb{E}_{\mathbf{E}}[V(\mathbf{E}_{\hat{\tau}_{n+1}^+}) \mid \tau_{n+1} > h] \cdot \mathbb{P}_{\mathbf{E}}[\tau_{n+1} > h] + \mathbb{E}_{\mathbf{E}}[V(\mathbf{E}_{\hat{\tau}_{n+1}^+}) \mid \tau_{n+1} \leq h] \cdot \mathbb{P}_{\mathbf{E}}[\tau_{n+1} \leq h] \\ &\leq \mathbb{E}_{\mathbf{E}}[V(\mathbf{E}_{\hat{\tau}_n^+}) \mid \tau_{n+1} > h] \cdot \mathbb{P}_{\mathbf{E}}[\tau_{n+1} > h] \\ &\quad + (\mathbb{E}_{\mathbf{E}}[V(\mathbf{E}_{\hat{\tau}_n^+}) \mid \tau_{n+1} \leq h] + CM') \cdot \mathbb{P}_{\mathbf{E}}[\tau_{n+1} \leq h] \\ &\leq \mathbb{E}_{\mathbf{E}}[V(\mathbf{E}_{\hat{\tau}_n^+})] + CM'(1 - e^{-hNK})^{n+1}. \end{aligned}$$

Adding up from $n = 1$ to ∞ , this gives

$$P^h V(\mathbf{E}) \leq \mathbb{E}_{\mathbf{E}}[V(\mathbf{E}_{\hat{\tau}_1^+})] + \frac{CM'}{e^{-hNK}}.$$

Let $h > 0$ be small enough so that

$$\mathbb{P}_{\mathbf{E}}[\tau_1 \leq h] = 1 - e^{-h\mathcal{R}} > \frac{h}{2}\mathcal{R}.$$

This is the only condition we impose on h . Such an h exists because of the bound $\mathcal{R} \leq NK$.

Notice that the energy exchange at τ_1 is independent of τ_1 . We have, by Lemma 4.6,

$$\begin{aligned} \mathbb{E}_{\mathbf{E}}[V(\mathbf{E}_{\hat{\tau}_1^+})] &= \mathbb{E}_{\mathbf{E}}[V(\mathbf{E}_{\tau_1^+}) | \tau_1 \leq h] \cdot \mathbb{P}_{\mathbf{E}}[\tau_1 \leq h] + V(\mathbf{E}) \cdot \mathbb{P}_{\mathbf{E}}[\tau_1 > h] \\ &\leq \left(V(\mathbf{E}) - \frac{C_1}{\mathcal{R}} V(\mathbf{E})^\alpha \right) \cdot \mathbb{P}_{\mathbf{E}}[\tau_1 \leq h] + V(\mathbf{E}) \cdot \mathbb{P}_{\mathbf{E}}[\tau_1 > h] \\ &\leq V(\mathbf{E}) - C_1 \frac{h}{2} V(\mathbf{E})^\alpha. \end{aligned}$$

This gives

$$P^h V(\mathbf{E}) \leq V(\mathbf{E}) - C_1 \frac{h}{2} V(\mathbf{E})^\alpha + CM' e^{hNK}$$

for any $\mathbf{E} \in B_0$.

To complete the proof of Theorem 4.1, it suffices to replace M' by a large enough number $M_0 > 1$ so that for $\mathbf{E} \in \{V(\mathbf{E}) > M_0\}$, the constant $CM' e^{hNK}$ is absorbed into $c_0 V(\mathbf{E})^\alpha$ for $c_0 := C_1 \frac{h}{4}$. □

The same calculation in the proof of Theorem 4.1 also yields

Corollary 4.7.

$$\sup_{\{\mathbf{E}: V(\mathbf{E}) \leq \hat{M}\}} P^h V(\mathbf{E}) \leq \hat{M} + CM' e^{hNK} < \infty.$$

for any $\hat{M} > 0$.

Proof. The calculation in (4.4) gives

$$\mathbb{E}_{\mathbf{E}}[V(\mathbf{E}_{\hat{\tau}_{n+1}^+})] \leq \mathbb{E}_{\mathbf{E}}[V(\mathbf{E}_{\hat{\tau}_n^+})] + CM'(1 - e^{-hNK})^{n+1}.$$

Adding up from $n = 0$ to ∞ , this gives

$$P^h V(\mathbf{E}) \leq V(\mathbf{E}) + CM' e^{hNK} \leq \hat{M} + CM' e^{hNK}.$$

□

5. EXCURSION TIME FOR THE INDUCED CHAIN

For sufficiently small given parameters $h > 0$ and $\eta > 0$, let M_0 be the constant defined in Theorem 4.1 and $B := \{\mathbf{E} | V(\mathbf{E}) > M_0\}$. Define $G = \mathbb{R}_+^N \setminus B$. The previous section gives bounds of the excursion time in B . As introduced in Section 3.2, now one should work on the G -induced chain. G is not a compact set as the energy at each site can be arbitrarily high. A common way to show the tightness of a Markov process on non-compact state space is to construct a Lyapunov function, as what we do in this section for the G -induced chain.

Let $h > 0$ be the given size of a time step (defined in Theorem 4.1). Process

$$\mathbf{E}_n = \mathbf{E}_{nh}, \quad n \geq 0$$

is said to be the time- h sampling chain of $\mathbf{E}_t$.

Let $0 = T_0 < T_1 < T_2 < \dots$ be stopping times such that

$$T_1 = \inf_{k>0} \{\mathbf{E}_k \in G\}$$

and

$$T_{n+1} = \inf_{k>T_n} \{\mathbf{E}_k \in G\}$$

for $n = 1, 2, \dots$. We define

$$\hat{\mathbf{E}}_n = \mathbf{E}_{T_n}$$

for $n = 1, 2, \dots$ as the G -induced chain. It is easy to see that $\hat{\mathbf{E}}_n$ is also a Markov chain. We denote the kernel of $\hat{\mathbf{E}}_n$ by $\hat{P}$.

As in [22], a natural Lyapunov function is the total energy in the system. Let

$$W(\mathbf{E}) = \sum_{n=1}^N E_n.$$

The main theorem of this section reads:

Theorem 5.1. *There exist constants $M_1 > 1$ and $\delta > 0$, such that*

$$\hat{P}W(\mathbf{E}) \leq (1 - \delta)W(\mathbf{E})$$

for every $\mathbf{E} \in G$ with $W(\mathbf{E}) > M_1$.

To prove Theorem 5.1, the first task is to bound the length of the time step for $\hat{\mathbf{E}}_n$. Such estimate follows from Theorem 4.1 and Theorem 3.12 immediately.

Proposition 5.2. *There exists constant C_5 such that*

$$\mathbb{E}[(T_{n+1} - T_n)^{\hat{\alpha}} | \hat{\mathbf{E}}_n] \leq C_5 \hat{V}(\hat{\mathbf{E}}_n)$$

for any $\hat{\mathbf{E}}_n$ and any $n \geq 0$, where $\hat{\alpha} = (1 - \alpha)^{-1} = 2 - 2\eta$ for the constant α given in Theorem 4.1, and $\hat{V}(\mathbf{E}) = \max\{V(\mathbf{E}), 1\}$. In particular, if $n \geq 1$, then

$$\mathbb{E}[(T_{n+1} - T_n)^{\hat{\alpha}} | \hat{\mathbf{E}}_n] \leq C_5 M_0,$$

where M_0 is as in Theorem 4.1.

Proof. By the definition of $\hat{\mathbf{E}}_n$, $\hat{\mathbf{E}}_n = \mathbf{E}_{T_n}$ is the energy configuration at the stopping time T_n . Notice that

$$\mathbb{E}[(T_{n+1} - T_n)^{\hat{\alpha}} | \hat{\mathbf{E}}_n] \leq 2 \cdot \mathbb{E}_{\mathbf{E}_{T_n}} \left[\sum_{k=0}^{\tau_G} (k+1)^{\hat{\alpha}-1} \right]$$

for the constant α we use. The proposition follows immediately by applying Theorem 3.12 to $\mathbf{E}_n$. Since $M_0 > 1$, let

$$\hat{V}(\mathbf{E}) = \max\{1, V(\mathbf{E})\}.$$

Then since $\hat{V} \leq V + 1$, it follows from Theorem 4.1 and Corollary 4.7 that

$$\hat{P}\hat{V}(\mathbf{E}) - \hat{V}(\mathbf{E}) \leq -c_0\hat{V}^\alpha(\mathbf{E}) + (1 + M_0 + CM'e^{hNK})\mathbf{1}_G.$$

The proposition then follows from Theorem 3.12. $\square$

The following definitions regarding the energy flux in the system are necessary for the proof of Theorem 5.1. Let $t_1, t_2, \dots$ be the times at which either clock 1 or clock $N + 1$ rings. The energy in-flow and out-flow on $[0, T)$ are denoted by

$$F_I([0, T)) = \sum_{0 \leq t_i < T} (W(\mathbf{E}_{t_i^+}) - W(\mathbf{E}_{t_i}))^+$$

and

$$F_O([0, T)) = \sum_{0 \leq t_i < T} (W(\mathbf{E}_{t_i}) - W(\mathbf{E}_{t_i^+}))^+,$$

respectively. Next we need to estimate the energy flux with respect to $\hat{\mathbf{E}}$.

Lemma 5.3. *There exists a constant C_2 such that*

$$\mathbb{E}_{\mathbf{E}}[F_I([0, T))] \leq C_2 T$$

for any $\mathbf{E} \in \mathbb{R}_+^N$ and any $T > 0$. In addition

$$X_n := F_I([0, nh)) - C_2 hn$$

is a super-martingale relative to $\mathcal{F}_n$, where $\mathcal{F}_n$ is the filtration generated by $\mathbf{E}_n$.

Proof. It is easy to see that for any $\mathbf{E}(t) \in \mathbb{R}_+^N$ we have

$$\begin{aligned} F_I([t, t + dt)) &\leq \sqrt{T_L} \left\{ \int_0^\infty \int_0^1 \frac{1}{T_L} p(x + E_1(t)) e^{-x/T_L} dx dp - E_1 \right\}^+ dt \\ &\quad + \sqrt{T_R} \left\{ \int_0^\infty \int_0^1 \frac{1}{T_R} p(x + E_N(t)) e^{-x/T_R} dx dp - E_N \right\}^+ dt \\ &\leq \sqrt{T_L} \left\{ \int_0^\infty \int_0^1 \frac{1}{T_L} p x e^{-x/T_L} dx dp \right\} dt + \sqrt{T_R} \left\{ \int_0^\infty \int_0^1 \frac{1}{T_R} p x e^{-x/T_R} dx dp \right\} dt \\ &= \frac{1}{2} (T_L^{3/2} + T_R^{3/2}) dt := C_2 dt. \end{aligned}$$

This estimate is independent of $\mathbf{E}(t)$. Therefore,

$$\mathbb{E}_{\mathbf{E}}[F_I([0, T))] \leq C_2 T$$

for any $\mathbf{E}$ and T . In addition,

$$\mathbb{E}[X_{n+1} - X_n \mid \mathcal{F}_n] = \mathbb{E}[F_I([nh, (n+1)h)) \mid \mathcal{F}_n] - C_2 h \leq 0$$

for any $\mathbf{E}_n$. This completes the proof. $\square$

By the definition of T_1 , T_1 is a stopping time relative to $\mathcal{F}_n$. We have

Proposition 5.4.

$$\mathbb{E}_{\mathbf{E}}[F_I([0, T_1))] \leq C_2 h \mathbb{E}_{\mathbf{E}}[T_1],$$

where C_2 is the constant in Lemma 5.3.

Proof. By Proposition 5.2, $\mathbb{E}_{\mathbf{E}}[T_1] < \infty$ for any $\mathbf{E} \in \mathbb{R}_+^N$. In addition, by Lemma 5.3,

$$\mathbb{E}_{\mathbf{E}}[|X_{n+1} - X_n| \mid \mathcal{F}_n] \leq 2C_2h < \infty,$$

where the super-martingale X_n is defined in Lemma 5.3. It then follows from the optional stopping theorem that

$$\mathbb{E}_{\mathbf{E}}[X_{T_1}] \leq \mathbb{E}_{\mathbf{E}}[X_0] = 0.$$

Therefore

$$\mathbb{E}_{\mathbf{E}}[F_I([0, T_1))] \leq C_2h\mathbb{E}_{\mathbf{E}}[T_1].$$

□

The following estimate about the energy influx with respect to the G -induced chain follows easily.

Lemma 5.5. *There exist constants $C_3 < \infty$ and $C'_3 < \infty$ such that*

$$\mathbb{E}_{\mathbf{E}}[F_I([0, T_1))] \leq C_3$$

if $\mathbf{E} \in G$ and

$$\mathbb{E}_{\mathbf{E}}[F_I([0, T_1))] \leq C'_3\hat{V}(\mathbf{E}).$$

if $\mathbf{E} \notin G$.

Proof. By Proposition 5.2,

$$\mathbb{E}_{\mathbf{E}}[T_1] \leq \mathbb{E}_{\mathbf{E}}[(T_1)^{\hat{\alpha}}] \leq C_5\hat{V}(\mathbf{E}).$$

It then follows from Proposition 5.4 that

$$\mathbb{E}_{\mathbf{E}}[F_I([0, T_1))] \leq C_2h \cdot C_5\hat{V}(\mathbf{E}).$$

Let

$$C_3 := C_2h \cdot C_5M_0.$$

If $\mathbf{E}_0 \in G$, the lemma follows immediately. If $\mathbf{E}_0 \notin G$, by letting

$$C'_3 = C_2hC_5,$$

we will have

$$\mathbb{E}_{\mathbf{E}}[F_I([0, T_1))] \leq C'_3\hat{V}(\mathbf{E}).$$

This completes the proof.

□

The following lemma controls the out flow of the energy.

Lemma 5.6. *Assume $\mathbf{E} \in G$. There exist constants $\sigma, M'' > 0$, such that*

$$\mathbb{E}_{\mathbf{E}}[F_O([0, T_1))] \geq \sigma W(\mathbf{E})$$

whenever $W(\mathbf{E}) > M''$.

Proof. Since

$$\mathbb{E}_{\mathbf{E}}[F_O([0, T_1))] \geq \mathbb{E}_{\mathbf{E}}[F_O([0, h))],$$

it is sufficient to construct an event within the time interval $[0, h)$ such that whenever $\mathbf{E} \in G$, a certain proportion of energy can be dumped out of the system.

Let E_{n_1} be the site that holds the largest amount of energy. Let $\mathcal{E}$ denote the following event

- Clocks $n_1, n_1 - 1, \dots, 2$, and 1 ring in the time interval $[0, \frac{h}{n_1}), [\frac{h}{n_1}, \frac{2h}{n_1}), \dots, [\frac{(n_1-1)h}{n_1}, h)$, respectively.
- At the i -th ring, E_{n_1-i+1} gives at least half of its energy to E_{n_1-i} for $i = 1, \dots, n_1 - 1$.
- At the n_1 -th ring, E_1 dumps $1/3$ of its energy to the left heat bath.
- Besides what described above, all other clocks do not ring during the time period $[0, h)$.

If $W(\mathbf{E}) \geq 3N \cdot 2^{N-1}T_L$, we have $E_{n_1}(0) \geq 3 \cdot 2^{N-1} \cdot T_L$ and $E_1(\frac{(n_1-1)h}{n_1}) > 3T_L$ conditioning with event $\mathcal{E}$. Note that all clock rates are bounded above by K . In addition, the probability that E_{n_1-i+1} gives at least half of its energy to E_{n_1-i} is at least 2^{-N} .

Therefore, it is a straightforward exercise to check that there exists a constant $c_0 > 0$ such that

$$\mathbb{P}[\mathcal{E}] \geq c_0 > 0$$

for every $\mathbf{E} \in G$. The proof is completed by letting

$$\sigma = \frac{c_0}{3} \cdot 2^{-(N-1)}$$

and

$$M'' = 3N \cdot 2^{N-1}T_L.$$

□

Proof. Proof of Theorem 5.1.

By the definition of $W(\mathbf{E})$, we have

$$\hat{P}W(\mathbf{E}) = \mathbb{E}_{\mathbf{E}}[F_I([0, T_1))] - \mathbb{E}_{\mathbf{E}}[F_O([0, T_1))].$$

Since $\mathbf{E} \in G$, by Lemma 5.5, we have $\mathbb{E}_{\mathbf{E}}[F_I([0, T_1))] \leq C_3$. On the other hand, by Lemma 5.6, if $W(\mathbf{E}) > M''$, we have

$$\mathbb{E}_{\mathbf{E}}[F_O([0, T_1))] \geq \sigma W(\mathbf{E}).$$

Therefore, let $\delta = \frac{1}{2}\sigma$ and $M_1 = \max\{2C_3/\sigma, 1\}$, we have

$$\hat{P}W(\mathbf{E}) \leq (1 - \delta)W(\mathbf{E}).$$

This completes the proof.

□

6. PROOF OF THE THEOREMS

6.1. Existence of uniform reference set $\mathfrak{C}$. Define

$$\mathfrak{C} = \{\mathbf{E} \mid V(\mathbf{E}) \leq M_0, W(\mathbf{E}) \leq M_1\}.$$

The aim of this subsection is to prove that $\mathfrak{C}$ is a uniform reference set. This follows immediate from the theorem below.

Theorem 6.1. *For any $t > 0$, there exists a constant $\eta > 0$ such that*

$$P^t(\mathbf{E}, \cdot) > \eta U_{\mathfrak{C}}(\cdot) \quad \text{for all } \mathbf{E} \in \mathfrak{C},$$

where $U_{\mathfrak{C}}$ is the uniform probability measure on $\mathfrak{C}$.

Proof. Let $e = \inf_{\mathbf{E}=(E_1, \dots, E_N) \in K} \min(E_1, \dots, E_N)$. Clearly $e > 0$.

Then we cover $\mathfrak{C}$ by finite many disks $D = D(\bar{\mathbf{E}}, \xi) = \{\mathbf{E} \mid |\mathbf{E} - \bar{\mathbf{E}}| \leq \xi\}$ for $\xi < e/2$. It is sufficient to show that for any $t > 0$, there exists $\eta > 0$ independent of $\bar{\mathbf{E}} \in K$, such that for any $\mathbf{E} \in K$, $P^t(\mathbf{E}, \cdot) \geq \eta U_D(\cdot)$ for all D in this cover.

Let $\mathbf{E}$ and D be fixed. We prescribe the following sequence of events.

- (i) On the time intervals $[\frac{(i-1)t}{2N}, \frac{it}{2N})$ for each $i = 1 \sim N-1$, site i exchanges energy with site $i+1$. After the energy exchange, the remaining energy at site i is between $e/2$ and e . Other clocks do not ring during this time period.
- (ii) On the time interval $[\frac{(N-1)t}{2N}, \frac{1}{2})$, site N exchanges energy with the right heat bath. After the energy exchange, E_N is greater than $\sup_{\bar{\mathbf{E}} \in D} \sum_{i=1}^N (\bar{E}_i + \xi)$. Other clocks do not ring during this time period.
- (iii) On the time intervals $[\frac{(N+i-1)t}{2N}, \frac{(N+i)t}{2N})$ for each $i = 1 \sim N$, site $N-i$ exchanges energy with site $N+1-i$. (Note that site 0 is the left heat bath.) After each energy exchange, the energy left at site $N+1-i$ is uniformly distributed in $E_{N+1-i} \in [\bar{E}_{N+1-i} - \xi, \bar{E}_{N+1-i} + \xi]$. Other clocks do not ring during this time period.

It is then easy to check that the event above occurs with probability at least η , where $\eta > 0$ is independent of $\mathbf{E}$ provided $\mathbf{E} \in \mathfrak{C}$. $\square$

Since $P^t(\mathbf{E}, \cdot)$ now has positive density everywhere in $\mathbb{R}_+^N$, the strong aperiodicity and irreducibility of $\mathbf{E}_n$ follows immediately.

Corollary 6.2. $\mathbf{E}_n$ is a strongly aperiodic Markov chain.

Proof. By theorem 6.1, $\mathfrak{C}$ is a uniform reference set. In addition $U_{\mathfrak{C}}(\mathfrak{C}) > 0$. The strong aperiodicity follows from its definition. $\square$

Therefore $\mathbf{E}_n$ is aperiodic.

Corollary 6.3. $\mathbf{E}_n$ is λ -irreducible, where λ is the Lebesgue measure on $\mathbb{R}_+^N$.

Proof. Let $A \subset \mathbb{R}_+^N$ be a set with strictly positive Lebesgue measure. Then there exists a set O that has the form

$$O = \{\mathbf{E} \mid 0 < c \leq E_i \leq C < \infty, i = 1 \sim N\}$$

such that $\lambda(O \cap A) > 0$.

For any $\mathbf{E}_0 \in \mathbb{R}_+^N$ and the time step h as in Theorem 4.1, same construction as in Theorem 6.1 implies that $P^h(\mathbf{E}_0, \cdot) > \eta U_O(\cdot)$ for some $\eta > 0$. Therefore, when ϵ is sufficiently small, $P^h(\mathbf{E}_0, A) > \eta U_O(A) > 0$. $\square$

6.2. Absolute continuity of invariant measure. This subsection aims to prove the absolute continuity of the invariant probability with respect to the Lebesgue measure. For the sake of simplicity, we denote the Lebesgue measure on $\mathbb{R}_+^N$ by λ .

Proposition 6.4. *If π is an invariant measure of $\mathbf{E}_t$, then π is absolutely continuous with respect to λ with a strictly positive density.*

The proof is similar to that of Proposition 6.1 of [21]. For $\mathbf{E} \in \mathbb{R}_+^N$ and $t > 0$, we have decomposition

$$P^t(\mathbf{E}, \cdot) = \nu_\perp + \nu_{abs},$$

where ν_{abs} and $\nu_\perp$ are absolutely continuous and singular component with respect to λ , respectively. We need to show that an absolutely continuous component cannot revert back to singularity as time evolves.

Lemma 6.5. *For any probability measure $\mu \ll \lambda$, $\mu P^t \ll \lambda$ for any $t > 0$.*

Proof. This proof is similar to that of Lemma 6.3 of [21]. We include it here for the sake of completeness of this paper.

Let $t > 0$ be fixed. We define $l = (c_1, \dots, c_n)$ be the sequence of energy exchanges, where $c_i = k$ means site k exchanges energy with site $k + 1$. (As before, heat baths are sites 0 and $N + 1$.) Let $S(l)$ be the event that energy exchanges $(c_1, \dots, c_n)$ occur during the time period $[0, t)$ in the order specified, and no other energy exchanges occur. If zero (resp. infinite many) energy exchange occurs on $[0, t)$, we denote the corresponding event by $S(\emptyset)$ (resp. $S(\infty)$). Obviously $\mathbb{P}_{\mathbf{E}}[S(\infty)] = 0$.

For $S = S(l)$ or $S(\emptyset)$, we define the conditional Markov operator

$$(\mu P_S)(A) := \int_{\mathbb{R}_+^N} \mathbb{P}[\mathbf{E}_t \in A \mid \mathbf{E}_0 = \mathbf{E} \mid S] \mu(d\mathbf{E})$$

and the measure

$$\frac{d\mu_Q}{d\lambda}(\mathbf{E}) := \mathbb{P}_{\mathbf{E}}[S(Q)] \frac{d\mu}{d\lambda}(\mathbf{E})$$

for $Q = l$ or $\emptyset$. Then by the law of total probability,

$$\mu P^t = \mu_\emptyset + \sum_l \mu_l P_{S(l)}.$$

Therefore, it is sufficient to show that each term above is absolutely continuous with respect to λ . Since for each $l = (c_1, \dots, c_n)$ we have the decomposition

$$P_{S(l)} = P_{S(c_n)} \cdots P_{S(c_1)}.$$

Hence the proof is reduced to proving the absolute continuity of μP_{c_i} for each $i = 0 \sim N + 1$, which is a straightforward exercise. Let ξ and $\hat{\xi}_k$ be the density of μ and

$\mu P_{S(c_k)}$, respectively. Then if $0 < k < N$, we have

$$\begin{aligned} & \hat{\xi}_k(E_1, E_2, \dots, E_N) \\ &= \int_0^1 \xi(E_1, \dots, E_{k-1}, p(E_k + E_{k+1}), (1-p)(E_k + E_{k+1}), E_{k+2}, \dots, E_N) dp. \end{aligned}$$

For $k = 0$ and N , we have

$$\begin{aligned} & \hat{\xi}_0(E_1, E_2, \dots, E_N) \\ &= \int_0^\infty \int_{(E_1 - \hat{E}_1)^+}^\infty \xi(\hat{E}_1, E_2, \dots, E_N) \frac{1}{\hat{E}_1 + E} \frac{1}{T_L} e^{-E/T_L} dE d\hat{E}_1 \end{aligned}$$

and

$$\begin{aligned} & \hat{\xi}_N(E_1, E_2, \dots, E_N) \\ &= \int_0^\infty \int_{(E_N - \hat{E}_N)^+}^\infty \xi(E_1, E_2, \dots, \hat{E}_N) \frac{1}{\hat{E}_N + E} \frac{1}{T_R} e^{-E/T_R} dE d\hat{E}_N. \end{aligned}$$

□

Proof of Proposition 6.4. Let $\pi = \pi_{abs} + \pi_{perp}$ be an invariant measure. Assume $\pi_{perp} \neq 0$. For $t > 0$, $\pi_{abs} P^t \ll \lambda$ by Lemma 6.4. By Theorem 6.1, for any $\mathbf{E} \in \mathfrak{C}$, $P^{t/2}(\mathbf{E}, \cdot)$ has a strictly positive density on $\mathfrak{C}$. Since $\mathfrak{C}$ is accessible within finite many energy exchanges, $P^{t/2}(\mathbf{E}, \mathfrak{C}) > 0$ for all $\mathbf{E} \in \mathbb{R}_+^N$. Hence $P^t(\mathbf{E}, \cdot)$ has a strictly positive density on $\mathfrak{C}$ for all $\mathbf{E} \in \mathbb{R}_+^N$. Therefore, $\pi_\perp P^t$ has an absolutely continuous component. The absolutely continuous component of πP^t is strictly large than that of π . This contradicts to the invariance of π .

□

6.3. Excursion time before entering $\mathfrak{C}$. Now we are ready to estimate the tail of $\tau_{\mathfrak{C}}$. Let $\hat{V} = \max\{1, V\}$ and $\hat{W} = \max\{1, W\}$.

Theorem 6.6. *For any $\epsilon > 0$, there exists a constant $C_6 < \infty$ such that*

$$\mathbb{P}_{\mathbf{E}}[\tau_{\mathfrak{C}} > n] \leq C_6(\hat{W}(\mathbf{E}) + \hat{V}(\mathbf{E}))n^{-(\hat{\alpha}-\epsilon)}$$

for any $\mathbf{E} \in \mathbb{R}_+^N$, where $\hat{\alpha} = 2 - 2\eta$.

Proof. Note that $\mathfrak{C} \subset G$, therefore one can define $\hat{\tau}_{\mathfrak{C}}$ as the first passage time to $\mathfrak{C}$ for the induced chain $\hat{\mathbf{E}}_n$.

Note that $M_0, M_1 > 1$. Apply Theorem 3.11 to $\hat{W} = \max\{1, W(\mathbf{E})\}$. It follows from Theorem 3.11 and Theorem 5.1 that for every $\mathbf{E} \in G$, there exists constants $C_7 > 0$ and $r > 1$ such that

$$\mathbb{E}_{\mathbf{E}}[r^{\hat{\tau}_{\mathfrak{C}}}] \leq \mathbb{E}_{\mathbf{E}}\left[\sum_{k=0}^{\tau_{\mathfrak{C}}} W(\hat{\mathbf{E}}_k) r^k\right] < \hat{W}(\mathbf{E}) C_7.$$

Applying Markov inequality to $r^{\hat{\tau}_{\mathfrak{C}}}$, we have

$$(6.1) \quad \mathbb{P}_{\mathbf{E}}[\hat{\tau}_{\mathfrak{C}} > n] < C_7 \hat{W}(\mathbf{E}) e^{-cn},$$

where $c = \log r$. For any given initial condition $\mathbf{E} \in \mathbb{R}_+^N$, we have

$$\begin{aligned}
& \mathbb{P}_{\mathbf{E}}[\hat{\tau}_{\mathfrak{C}} > n] \\
&= \int_{\mathbb{R}_+^N} \mathbb{P}[\hat{\tau}_{\mathfrak{C}} > n \mid \mathbf{E}_{T_1} = \tilde{E}] \mathbb{P}_{\mathbf{E}}[\mathbf{E}_{T_1} = d\tilde{E}] \\
&\leq \int_{\mathbb{R}^N} C_7 \hat{W}(\tilde{E}) e^{-cn} \mathbb{P}_{\mathbf{E}}[\mathbf{E}_{T_1} = \tilde{E}] \\
&= C_7 e^{-cn} \mathbb{E}_{\mathbf{E}}[\hat{W}(\mathbf{E}_{T_1})] \\
&\leq C_7 e^{-cn} \cdot (\hat{W}(\mathbf{E}) + \mathbb{E}_{\mathbf{E}}[F_I([0, T_1)])] \\
&\leq C_7 e^{-cn} \cdot (\hat{W}(\mathbf{E}) + \max\{C_3, C'_3 \hat{V}(\mathbf{E})\}) \\
&\leq C_7 e^{-cn} \cdot (\hat{W}(\mathbf{E}) + \max\{C_3, C'_3\} \hat{V}(\mathbf{E})),
\end{aligned}$$

where the third line follows from Equation (6.1) and the second to last line follows from Lemma 5.5, constants C_3 and C'_3 are as in Lemma 5.5.

By Proposition 5.2, we have

$$\mathbb{E}[(T_{n+1} - T_n)^{\hat{\alpha}} \mid \hat{\mathbf{E}}_n] \leq C_5 \hat{V}(\hat{\mathbf{E}}_n),$$

where C_5 is from Proposition 5.2. Without loss of generality, we let $C_5 \geq 1$.

Applying Markov inequality to $(T_{n+1} - T_n)^{\hat{\alpha}}$, we have

$$\mathbb{P}[T_{n+1} - T_n > k \mid \mathbf{E}_{T_n}] \leq C_5 \hat{V}(\mathbf{E}_{T_n}) k^{-\hat{\alpha}}.$$

If $n \geq 1$, since $\mathbf{E}_{T_n} = \hat{\mathbf{E}}_n \in G$, we have a uniform bound $\hat{V}(\mathbf{E}_0) \leq M_0$.

Therefore, assumptions of Theorem 3.8 are satisfied for $\xi = \hat{V}$ and $\eta = \hat{W}$. For each given $\epsilon > 0$, notice that $\hat{V} \geq 1$ and $\hat{W} \geq 1$, we have

$$\mathbb{P}_{\mathbf{E}}[\hat{\tau}_{\mathfrak{C}} > n] \leq c(C_5 \hat{V}(\mathbf{E}) + \hat{W}(\mathbf{E}) + \max\{C_3, C'_3\} \hat{V}(\mathbf{E})) n^{-(\hat{\alpha}-\epsilon)} \leq C_6 (\hat{W}(\mathbf{E}) + \hat{V}(\mathbf{E})) n^{-(\hat{\alpha}-\epsilon)},$$

where C_6 is a constant depending on ϵ and η . This completes the proof. $\square$

Proof of Theorem 1 and 2: We first prove Theorem 1 and 2 for $\mathbf{E}_n$.

It follows from Theorem 6.6 that

$$\mathbb{P}_{\mathbf{E}}[\tau_{\mathfrak{C}} > n] \leq C_6 (\hat{W}(\mathbf{E}) + \hat{V}(\mathbf{E})) n^{-(\hat{\alpha}-\epsilon)},$$

where C_6 is a constant that depends on both η and ϵ . Since $\mathbb{P}_{\mathbf{E}}[\tau_{\mathfrak{C}} > n]$ is at most 1, we have

$$\mathbb{P}_{\mathbf{E}}[\tau_{\mathfrak{C}} > n] \leq \max\{1, 2C_6\} (\hat{W}(\mathbf{E}) + \hat{V}(\mathbf{E})) (n+1)^{-(\hat{\alpha}-\epsilon)}.$$

Therefore, by Lemma 3.10, there exists a constant C_8 that depends on η and ϵ , such that

$$\mathbb{E}_{\mathbf{E}}[\tau_{\mathfrak{C}}^{\hat{\alpha}-2\epsilon}] \leq C_8 (\hat{W}(\mathbf{E}) + \hat{V}(\mathbf{E})).$$

Note that $\hat{\alpha} = 2 - 2\eta$. Let $\epsilon = \eta$ and $\eta = \frac{1}{4}\gamma$, we have

$$(6.2) \quad \mathbb{E}_{\mathbf{E}_0}[\tau_{\mathfrak{C}}^{2-\gamma}] \leq C_8 (\hat{W}(\mathbf{E}) + \hat{V}(\mathbf{E})).$$

By the compactness of $\mathfrak{C}$, it is easy to see that

$$\sup_{\mathbf{E} \in \mathfrak{C}} \mathbb{E}_{\mathbf{E}}[\tau_{\mathfrak{C}}^{2-\gamma}] < \infty.$$

Note that $\hat{W} \leq W + 1$ and $\hat{V} \leq V + 1$. Therefore, for any probability measure $\mu \in \mathcal{M}_{\gamma}$, $C_8(\hat{W}(\mathbf{E}) + \hat{V}(\mathbf{E}))$ is integrable and

$$(6.3) \quad \mathbb{E}_{\mu}[\tau_{\mathfrak{C}}^{2-\gamma}] < \infty.$$

Theorem 1 for $\mathbf{E}_n$ is then proved by applying Theorem 3.4 to $\mu, \nu \in \mathcal{M}_{\gamma}$.

It follows from Corollary 6.2 and 6.3 that $\mathbf{E}_n$ is a strongly aperiodic irreducible Markov chain. The existence of an invariant measure π then follows from

$$\sup_{\mathbf{E} \in \mathfrak{C}} \mathbb{E}_{\mathbf{E}}[\tau_{\mathfrak{C}}] < \sup_{\mathbf{E} \in \mathfrak{C}} \mathbb{E}_{\mathbf{E}}[\tau_{\mathfrak{C}}^{2-\gamma}] < \infty$$

and Theorem 3.3. The absolute continuity of π comes from Proposition 6.4. The $n^{-(1-\gamma)}$ speed of convergence to π is given by the existence of π and Theorem 3.4.

It remains to prove the uniqueness of π . We prove the uniqueness of π for any $\mathbf{E}_t$ instead of $\mathbf{E}_n$.

Recall the proof of Proposition 6.4, for any $t > 0$ and $\mathbf{E} \in \mathbb{R}_+^N$, $P^t(\mathbf{E}, \cdot)$ has a strictly positive density on $\mathfrak{C}$. This implies that all $\mathbf{E} \in \mathbb{R}_+^N$ are in the same ergodic component. Hence $\mathbf{E}_t$ has at most one invariant probability measure, which must be π .

This completes the proof of Theorem 2 for $\mathbf{E}_n$.

Theorem 1 and 2 for $\mathbf{E}_t$. The last step is to pass results from $\mathbf{E}_n$ to $\mathbf{E}_t$. The contraction of Markov operator is easy to pass because we have

$$\|\mu P^t - \nu P^t\|_{\text{TV}} = \|(\mu P^{\lfloor \frac{t}{h} \rfloor h} - \nu P^{\lfloor \frac{t}{h} \rfloor h}) P^{(t - \lfloor \frac{t}{h} \rfloor h)}\|_{\text{TV}} \leq \|\mu P^{\lfloor \frac{t}{h} \rfloor h} - \nu P^{\lfloor \frac{t}{h} \rfloor h}\|_{\text{TV}}.$$

Remark: We believe Theorem 1 is close to optimal. Let $\mathfrak{C}$ be a uniform reference set. Then it is easy to see that there exists an $\epsilon > 0$ such that $B_{\epsilon} = \{\mathbf{E} \mid E_i < \epsilon \text{ for some } i = 1 \sim N\}$ is disjoint with $\mathfrak{C}$. Then it is easy to see that for any μ and ν that has uniformly positive density on B_{ϵ} , the probability that the initial distribution $\mathbf{E}_0 \sim \mu$ (and $\mathbf{E}_0 \sim \nu$) satisfies $E_i(0) < t^{-2}$ is $\sim O(t^{-2})$ for $t \gg 1$. When $E_i(0) < t^{-2}$, the probability that no energy exchange occurs between site $i - 1$ and i , or between site i and $i + 1$, before time t is $O(1)$. Therefore, we have the lower bound of tails $\mathbb{P}_{\mu}[\tau_{\mathfrak{C}} > t] \geq O(1) \cdot t^{-2}$ (and $\mathbb{P}_{\nu}[\tau_{\mathfrak{C}} > t] \geq O(1) \cdot t^{-2}$). This implies the coupling time T has the tail

$$\mathbb{P}_{\mu, \nu}[T > t] \geq O(1) \cdot t^{-2}.$$

It remains to show that π , the invariant probability measure of $\mathbf{E}_n$, is invariant for any $\mathbf{E}_t$, $t > 0$.

Lemma 6.7. $\pi P^t = \pi$ for any $t > 0$.

Proof. Note that the argument in Section 4 and 5 works for all sufficiently small time steps. Let h be the time step we have chosen for $\mathbf{E}_n$. For any $r < h$, P^r also admits an invariant measure π_r . It is then sufficient to show that $\pi_r = \pi$ because any t can be written as

$$t = \lfloor \frac{t}{h} \rfloor \cdot h + r$$

for some $r < h$.

In addition, we have the “continuity at zero”.

$$\|\pi_r P^\delta - \pi_r\|_{TV} \rightarrow 0 \quad \text{as } \delta \rightarrow 0$$

because all clock rates are less than K .

Without loss of generality, assume $r/h \notin \mathbb{Q}$. By the density of orbits in irrational rotations, there exists sequences $i_n, j_n \in \mathbb{Z}^+$, such that $\delta_n := h - \frac{i_n}{j_n}r \rightarrow 0$ from right. Then

$$\pi_r P^h = \pi_r P^{\frac{i_n}{j_n}r} P^{\delta_n} = \pi_r P^{\delta_n} \rightarrow \pi_r$$

by the “continuity at zero”.

Hence π_r is invariant with respect to P^h . By uniqueness, $\pi_r = \pi$.

□

Therefore, Theorem 1 and 2 also hold for $\mathbf{E}_t$.

Proof of Theorem 3. As a direct consequence of Theorem 1, we have

$$\begin{aligned} & \left| \int (P^t \zeta)(\mathbf{E}) \xi(\mathbf{E}) \mu(d\mathbf{E}) - \int (P^t \zeta)(\mathbf{E}) \mu(d\mathbf{E}) \int \xi(\mathbf{E}) \mu(d\mathbf{E}) \right| \\ &= \left| \int \xi(\mathbf{E}) \left((P^t \zeta)(\mathbf{E}) - \int (P^t \zeta)(\mathbf{Z}) \mu(d\mathbf{Z}) \right) \mu(d\mathbf{E}) \right| \\ &\leq \|\xi\|_{L^\infty} \|\zeta\|_{L^\infty} \int \|\delta_{\mathbf{E}} P^t - \mu P^t\|_{TV} \mu(d\mathbf{E}). \end{aligned}$$

It then follows from Corollary 3.7 and equation (6.2) that

$$\|\delta_{\mathbf{E}} P^t - \mu P^t\|_{TV} \leq C(\hat{W}(\mathbf{E}) + \hat{V}(\mathbf{E}) + C_\mu)(\lfloor t \rfloor)^{\gamma-2}$$

for some $C, C_\mu < \infty$ that is independent of $\mathbf{E}$. Since $\hat{W}(\mathbf{E}) + \hat{V}(\mathbf{E})$ is μ -integrable, we have

$$\left| \int (P^t \zeta)(\mathbf{E}) \xi(\mathbf{E}) \mu(d\mathbf{E}) - \int (P^t \zeta)(\mathbf{E}) \mu(d\mathbf{E}) \int \xi(\mathbf{E}) \mu(d\mathbf{E}) \right| \leq O(1) \cdot t^{\gamma-2}$$

ACKNOWLEDGEMENT

The author would like to thank Lai-Sang Young, Jonathan Mattingly, and Martin Hairer for many enlightening discussions.

REFERENCES

- [1] Vlad Stefan Barbu and Nikolaos Limnios, *Semi-markov chains and hidden semi-markov models toward applications: their use in reliability and dna analysis*, vol. 191, Springer Science & Business Media, 2009.
- [2] Cédric Bernardin and Stefano Olla, *Fouriers law for a microscopic model of heat conduction*, Journal of Statistical Physics **121** (2005), no. 3, 271–289.
- [3] Leonid Bunimovich, Carlangelo Liverani, Alessandro Pellegrinotti, and Yurii Suhov, *Ergodic systems of n balls in a billiard table*, Communications in mathematical physics **146** (1992), no. 2, 357–396.
- [4] Nikolai Chernov, *Decay of correlations and dispersing billiards*, Journal of Statistical Physics **94** (1999), no. 3-4, 513–556.
- [5] Nikolai Chernov and Roberto Markarian, *Chaotic billiards*, no. 127, American Mathematical Soc., 2006.
- [6] Nikolai Chernov and Lai-Sang Young, *Decay of correlations for lorentz gases and hard balls*, Hard ball systems and the Lorentz gas, Springer, 2000, pp. 89–120.
- [7] Noé Cuneo and J-P Eckmann, *Non-equilibrium steady states for chains of four rotors*, Communications in Mathematical Physics (2016), 1–37.
- [8] Noé Cuneo, Jean-Pierre Eckmann, and Christophe Poquet, *Non-equilibrium steady state and subgeometric ergodicity for a chain of three coupled rotors*, Nonlinearity **28** (2015), no. 7, 2397.
- [9] Dmitry Dolgopyat and Carlangelo Liverani, *Energy transfer in a fast-slow hamiltonian system*, Communications in Mathematical Physics **308** (2011), no. 1, 201–225.
- [10] J-P Eckmann, C-A Pillet, and Luc Rey-Bellet, *Non-equilibrium statistical mechanics of anharmonic chains coupled to two heat baths at different temperatures*, Communications in Mathematical Physics **201** (1999), no. 3, 657–697.
- [11] J-P Eckmann and L-S Young, *Nonequilibrium energy profiles for a class of 1-d models*, Communications in Mathematical Physics **262** (2006), no. 1, 237–267.
- [12] Pierre Gaspard and Thomas Gilbert, *Heat conduction and fourier’s law in a class of many particle dispersing billiards*, New Journal of Physics **10** (2008), no. 10, 103004.
- [13] ———, *Heat conduction and fouriers law by consecutive local mixing and thermalization*, Physical review letters **101** (2008), no. 2, 020601.
- [14] Alexander Grigo, Konstantin Khanin, and Domokos Szasz, *Mixing rates of particle systems with energy exchange*, Nonlinearity **25** (2012), no. 8, 2349.
- [15] Martin Hairer, *Convergence of markov processes*, lecture notes (2010).
- [16] Søren F Jarner, Gareth O Roberts, et al., *Polynomial convergence rates of markov chains*, The Annals of Applied Probability **12** (2002), no. 1, 224–247.
- [17] C Kipnis, C Marchioro, and E Presutti, *Heat flow in an exactly solvable model*, Journal of Statistical Physics **27** (1982), no. 1, 65–74.
- [18] Raphaël Lefevere and Lorenzo Zambotti, *Hot scatterers and tracers for the transfer of heat in collisional dynamics*, Journal of Statistical Physics **139** (2010), no. 4, 686–713.
- [19] Yao Li, *On the stochastic behaviors of locally confined particle systems*, Chaos: An Interdisciplinary Journal of Nonlinear Science **25** (2015), no. 7, 073121.
- [20] Yao Li and Lai-Sang Young, *Existence of nonequilibrium steady state for a simple model of heat conduction*, Journal of Statistical Physics **152** (2013), no. 6, 1170–1193.
- [21] ———, *Nonequilibrium steady states for a class of particle systems*, Nonlinearity **27** (2014), no. 3, 607.
- [22] ———, *Polynomial convergence to equilibrium for a system of interacting particles*, Annals of Applied Probability (accepted).
- [23] Torngny Lindvall, *Lectures on the coupling method*, Courier Dover Publications, 2002.
- [24] Carlangelo Liverani and Stefano Olla, *Toward the fourier law for a weakly interacting anharmonic crystal*, Journal of the American Mathematical Society **25** (2012), no. 2, 555–583.

- [25] Sean P Meyn and Richard L Tweedie, *Stability of markovian processes iii: Foster-lyapunov criteria for continuous-time processes*, Advances in Applied Probability (1993), 518–548.
- [26] ———, *Markov chains and stochastic stability*, Cambridge University Press, 2009.
- [27] Esa Nummelin, *A splitting technique for harris recurrent markov chains*, Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete **43** (1978), no. 4, 309–318.
- [28] Esa Nummelin and Pekka Tuominen, *The rate of convergence in orey’s theorem for harris recurrent markov chains with applications to renewal theory*, Stochastic Processes and Their Applications **15** (1983), no. 3, 295–311.
- [29] Luc Rey-Bellet, *Nonequilibrium statistical mechanics of open classical systems*, XIVTH International Congress on Mathematical Physics, 2003, pp. 447–454.
- [30] Luc Rey-Bellet and Lawrence E Thomas, *Asymptotic behavior of thermal nonequilibrium steady states for a driven chain of anharmonic oscillators*, Communications in Mathematical Physics **215** (2000), no. 1, 1–24.
- [31] ———, *Fluctuations of the entropy production in anharmonic chains*, Annales Henri Poincare, vol. 3, Springer, 2002, pp. 483–502.
- [32] David Ruelle, *A mechanical model for fouriers law of heat conduction*, Communications in Mathematical Physics **311** (2012), no. 3, 755–768.
- [33] Makiko Sasada, *Spectral gap for stochastic energy exchange model with non-uniformly positive rate function*, arXiv preprint arXiv:1305.4066 (2013).
- [34] Tatiana Yarmola, *Sub-exponential mixing of open systems with particle–disk interactions*, Journal of Statistical Physics (2013), 1–20.
- [35] ———, *Sub-exponential mixing of random billiards driven by thermostats*, Nonlinearity **26** (2013), no. 7, 1825.

YAO LI: DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MASSACHUSETTS
AMHERST, AMHERST, MA, 01003

E-mail address: yaoli@math.umass.edu